

## List of Tables

| Table No. |                                                                                                                                                                                                                             | Page No. |
|-----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|
| 1.        | List of the conserved residues with corresponding natural variants.                                                                                                                                                         | 26       |
| 2.        | 2A: Primers used for insertion of Restriction sites in pBTR1                                                                                                                                                                | 36       |
| 3.        | 2B: Primers used for creating the N-terminal deletants of yeast iso-1-cyt- <i>c</i>                                                                                                                                         | 36       |
| 4.        | Interactions of extra 5 N-terminal residues with other residues of the main chain                                                                                                                                           | 49       |
| 5.        | $\alpha$ -helical content of WT, $\Delta(-5/-2)$ and $\Delta(-5/-3)$ yeast iso-1-cyt- <i>c</i>                                                                                                                              | 54       |
| 6.        | Thermal denaturation parameters of WT, $\Delta(-5/-2)$ and $\Delta(-5/-3)$ yeast iso-1-cyt- <i>c</i> obtained from changes in $[\theta]_{222}$ and $\Delta\varepsilon_{405}$ nm in native buffer at pH 6.0.                 | 58       |
| 7.        | Thermodynamic parameters of WT, and $\Delta(-5/-2)$ and $\Delta(-5/-3)$ yeast iso-1-cyt- <i>c</i> in native buffer obtained from DSC measurements in native buffer at pH 6.0                                                | 60       |
| 8.        | Thermodynamic parameters associated with GdmCl-induced denaturation of WT, $\Delta(-5/-2)$ and $\Delta(-5/-3)$ yeast iso-1-cyt- <i>c</i> at pH 6.0 and 25 °C                                                                | 68       |
| 9.        | Thermodynamic parameters associated with the heat-induced denaturation equilibrium, N $\leftrightarrow$ D state of $\Delta(-5/-2)$ yeast iso-1-cyt- <i>c</i> in the presence of different concentrations of GdmCl at pH 6.0 | 71       |
| 10.       | Thermodynamic parameters associated with the heat-induced denaturation equilibrium, N $\leftrightarrow$ D state of $\Delta(-5/-3)$ yeast iso-1-cyt- <i>c</i> in the presence of different concentrations of GdmCl at pH 6.0 | 74       |
| 11.       | Thermodynamic parameters associated with urea-induced denaturation of $\Delta(-5/-2)$ and $\Delta(-5/-3)$ yeast iso-1-cyt- <i>c</i> at pH 6.0 and 25 $\pm$ 0.1 °C                                                           | 83       |
| 12.       | Thermodynamic parameters associated with the heat-induced denaturation equilibrium, N $\leftrightarrow$ D state of $\Delta(-5/-2)$ iso-1-cyt- <i>c</i> in the presence of different concentrations of urea at pH 6.0        | 86       |
| 13.       | Thermodynamic parameters associated with the heat-induced denaturation equilibrium, N $\leftrightarrow$ D state of $\Delta(-5/-3)$ iso-1-cyt- <i>c</i> in the presence of different concentrations of urea at pH 6.0        | 86       |
| 14.       | Thermodynamic parameters associated with LiCl-induced denaturation of $\Delta(-5/-2)$ and $\Delta(-5/-3)$ yeast iso-1-cyt- <i>c</i> at pH 6.0 and 25 °C.                                                                    | 106      |
| 15.       | Comparison of the different structural properties of different states of WT, $\Delta(-5/-2)$ and $\Delta(-5/-3)$ yeast iso-1-cyt- <i>c</i> at pH 6.0 and 25 °C.                                                             | 112      |