

Figure No.	List of Figures	Page No.
1	Levels of protein structure. The repeating unit $-C-O-N-C^{\alpha}-$ makes up the backbone of protein, and R_1 , R_2 , R_3 and R_4 represent the various side chains. Arrow denotes β -strand and coil represent α -helix.	2
2	The protein folding funnel diagram shows proteins fold into their native structures by minimizing their free energy.	7
3	Structural view of the four thermodynamic states of a protein in terms of secondary and tertiary structure. These states are unfolded; PMG, MG and native state.	9
4	Stabilizing and destabilizing forces on the folding of a protein.	14
5	Diagram showing the burial of hydrophobic moieties and formation of intramolecular H-bonds upon protein folding.	17
6	Potential energy diagram showing the interaction between a pair of neutral atoms or molecules.	21
7	A schematic diagram showing protein stabilization brought about by the action of osmolyte.	44
8	Structure of mammalian kidney, the outer most region is the renal cortex while as the inner region is the renal medulla.	55
9	Schematic illustration of the molecular mechanisms involved in the adaptation of mammalian cells to hypertonic stress.	58
10	Display of the side chain and peptide backbone unit transfer free energies.	61
11	The Transfer Model for predicting the stability of proteins in osmolyte solution.	63
12	Schematic representation for preferential interaction.	65
13	Effect of surface tension on protein folding and unfolding.	66
14	Geometrical illustration of the excluded volume effect.	68
15	Elution profile plot of mA_{280} (milliabsorbance at 280 nm) versus elution volume (ml) of weak anion exchanger Hi Trap DEAE FF.	80
16	Elution profile of gel filtration chromatography.	81
17	SDS-PAGE of peaks obtained during gel filtration chromatography.	81

18	Elution profile plot of mA280 (milli absorbance at 280 nm) versus elution volume (ml) of weak anion exchanger Hi Trap DEAE FF.	82
19	Elution profile of strong anion exchanger Mono Q.	83
20	SDS-PAGE of peaks obtained during weak anion exchange and strong anion exchange chromatography.	83
21	Absorption spectrum of CRT-DNA Complex.	85
22	Agarose gel electrophores of CRT showing the presence of DNA.	85
23	Elution profile of strong anion exchanger Mono Q column.	86
24	SDS-PAGE of peaks obtained during Mono-Q exchange chromatography of 100% ammonium sulphate precipitated protein.	87
25	Urea-induced denaturation of SSA monitored by change in $\Delta\epsilon_{287}$ at pH 7.4 and 25 °C.	91
26	Urea-induced denaturation of SSA monitored by change in $[\theta]_{222}$ at pH 7.4 and 25 °C.	93
27	Urea-induced denaturation of SSA monitored by change in F_{347} at pH 7.4 and 25 °C.	95
28	Renaturation of the urea-induced denatured SSA at pH 7.4 and 25 °C	97
29	The far-UV CD spectra of various states of SSA at pH 7.4 and 25 °C.	98
30	The near-UV CD spectra of various states of SSA at pH 7.4 and 25 °C.	99
31	Fluorescence emission spectra of ANS bound to the native SSA, urea-induced X state and urea-induced D state at pH 7.4 and 25 °C.	99
32	Urea-induced unfolding of SSA as followed by $[\theta]_{222}$ at pH 7.4, 25 °C in the absence and presence of sorbitol.	102
33	Urea-induced unfolding of SSA as followed by F_{347} at pH 7.4, 25 °C in the absence and presence of sorbitol.	103
34	Urea-induced unfolding of SSA as followed by $[\theta]_{222}$ at pH 7.4, 25 °C in the presence of myo-inositol and glycine betaine.	104
35	Urea-induced unfolding of SSA as followed by F_{347} at pH 7.4, 25 °C in the absence and presence of 0.2 M myo-inositol and 1.0 M glycine betaine.	105

36	Reversibility of urea-induced denaturation of SSA in the absence and presence of different concentrations of sorbitol at pH 7.4 and 25 °C.	107
37	Reversibility of urea-induced denaturation of SSA in the presence of different concentrations of myo-inositol at pH 7.4 and 25 °C.	108
38	Reversibility of urea-induced denaturation of SSA in the presence of different concentrations of glycine betaine at pH 7.4 and 25 °C	109
39	Reversibility of urea-induced denaturation of SSA monitored by intrinsic fluorescence in the absence of osmolyte.	110
40	GdmCl-induced denaturation of SSA monitored by change in $[\theta]_{222}$ at pH 7.4 and 25 °C.	113
41	Renaturation of the GdmCl-induced denatured SSA at pH 7.4 and 25 °C.	115
42	The far-UV CD spectra of various states of SSA at pH 7.4 and 25 °C: Native protein (curve 1); GdmCl-induced X- state (curve 2); and GdmCl-induced D state (curve 3).	115
43	The near-UV CD spectra of various states of SSA at pH 7.4 and 25 °C: Native protein (curve 1); GdmCl-induced X- state (curve 2); and GdmCl-induced D state (curve 3).	116
44	Fluorescence emission spectra of ANS bound to the native SSA (curve 1), GdmCl-induced X state (curve 2) and GdmCl-induced D state (curve 3) at pH 7.4 and 25 °C. The excitation wavelength was 380 nm.	117
45	GdmCl-induced unfolding of SSA as followed by $[\theta]_{222}$ at pH 7.4, 25 °C in the absence and presence of (A) sorbitol, (B) Myo-inositol and (C) glycine betaine.	119
46	Normalized transition curves (A) $N \leftrightarrow X$, $X \leftrightarrow D$ (in the absence of sorbitol) & $N \leftrightarrow D$ (in the presence of 1.0 M sorbitol) as function of $[GdmCl]$ and (B) plots of ΔG_I , ΔG_{II} (in the absence of sorbitol) and ΔG_D (in the presence of 1.0 M sorbitol) as function of $[GdmCl]$.	119
47	GdmCl-induced unfolding of SSA as followed by F_{347} at pH 7.4, 25 °C in the (A) absence and presence of 1.0 M sorbitol (B) in the absence and presence of 0.2 M Myo-inositol and (C) in the absence and presence of 1.0 M glycine betaine.	121

48	Reversibility of GdmCl-induced denaturation of SSA in the presence of different concentrations of sorbitol at pH 7.4 and 25 °C.	123
49	Reversibility of GdmCl-induced denaturation of SSA in the presence of different concentrations of myo-inositol at pH 7.4 and 25 °C.	124
50	Reversibility of GdmCl-induced denaturation of SSA in the presence of different concentrations of glycine betaine at pH 7.4 and 25 °C.	125
51	Reversibility of GdmCl-induced denaturation of SSA monitored by intrinsic fluorescence in the absence of osmolyte.	126
52	GdmCl-induced denaturation of SSA at different urea concentrations monitored at $[\theta]_{222}$ at pH 7.4 and 25 °C.	127
53	GdmCl-induced denaturation curves of SSA in the presence of different urea-glycine betaine mixtures [urea] + [glycine betaine] at pH 7.4 and 25 °C.	129
54	GdmCl-induced denaturation curves of SSA in the presence of different urea- myo-inositol mixtures [urea] + [myo-inositol] at pH 7.4 and 25 °C.	131
55	GdmCl-induced denaturation curves of SSA in the presence of different urea-sorbitol mixtures [urea] + [sorbitol] at pH 7.4 and 25 °C.	133
56	Urea and GdmCl denaturation of CRT-DNA complex.	135
57	f_D plots of CRT-DNA complex at different concentrations of (A) myo-inositol, (B) glycine betaine and (C) sorbitol.	137
58	GdmCl-induced denaturation curves of CRT-DNA complex in the presence and absence of different concentrations of sorbitol followed by changes at $[\theta]_{222}$ and F_{347} .	138
59	Normalized transition curves of CRT-DNA complex $N \leftrightarrow X$ and $X \leftrightarrow D$ in the absence of sorbitol (red) and presence of 0.4 M sorbitol (blue) and 0.8 M sorbitol (green).	139
60	GdmCl-induced denaturation curves of CRT-DNA complex in the absence and presence of different concentrations of myo-inositol followed by changes at $[\theta]_{222}$ and F_{347} .	140
61	Normalized transition curves of CRT-DNA complex $N \leftrightarrow X$ and $X \leftrightarrow D$ in the absence and presence of myo-inositol.	141
62	GdmCl-induced denaturation curves of CRT-DNA complex in the absence and presence of different concentrations of glycine betaine followed by changes at $[\theta]_{222}$ and F_{347} .	142

63	Normalized transition curves of CRT-DNA complex $N \leftrightarrow X$ and $X \leftrightarrow D$ in the absence and presence of glycine betaine.	143
64	Crystal structure of Sheep serum albumin.	146
65	X ray crystallography structure of calnexin which has high amino acid homology to CRT.	149
66	Urea-induced denaturation curves of SSA (A) and CRT-DNA complex (B) monitored at $[\theta]_{222} / CD$, mdeg against [urea].	152
67	GdmCl-induced denaturation curves of SSA (A) and CRT-DNA complex (B) monitored at $[\theta]_{222} / CD$, mdeg against [GdmCl].	154
68	Effect of co-solutes and their mixtures on the secondary structure of the native and denatured states of SSA.	157
69	Normalized transition curves of $[\theta]_{222}$ and F_{347} of SSA induced by urea in the presence of highest concentration of osmolytes at pH 7.4 and 25 °C.	163
70	Normalized transition curves of $[\theta]_{222}$ and F_{347} of SSA induced by GdmCl in the presence of highest concentration of osmolytes at pH 7.4 and 25 °C.	164
71	Effect of myo-inositol, glycine betaine, sorbitol, urea and their predicted mixtures on stability of SSA at pH 7.4 and 25 °C values.	167