



## *References*

- Abdel-Massih R, Elias Abdou E, Baydoun E and Ziad Daoud Z (2010) Antibacterial Activity of the extracts obtained from *Rosmarinus officinalis*, *Origanum majorana*, and *Trigonella foenum-graecum* on highly drug-resistant gram negative bacilli. *J Botany Article ID* 464087, 8 pages doi:10.1155/2010/464087.
- Acharya SN, Sirchamroen A, Basu S, Ooraikar B and Basu SK (2006) Improvement in the nutraceutical properties of fenugreek (*T. foenum-graecum*). *Songkanakarin J Sci Technol* 28: 1-9.
- Acharya SN, Thomas JE and Basu SK (2007) Fenugreek: an “old world” crop for the “new world”. *Biodiversity* 7: 27-30.
- Acharya SN, Thomas JE, Prasad R and Basu SK. Diseases of fenugreek (*Trigonella foenum-graecum*L.) and their control measures with special emphasis on fungal diseases. In Acharya A and Perella AE eds. Management of fungal plant pathogens. CAB International, 2010, pp 245-261.
- Adesokan AA, Yakubu MT, Owoyele BV, Akanji MA, Soladoye A and Lawal OK (2008) Effect of administration of aqueous and ethanolic extracts of *Enantia chlorantha* stem bark on brewer's yeast-induced pyresis in rats. *Afr J Biochem Res* 2: 165-169.
- Agarwal K and Gupta PK (1983) Cytological studies in genus *Trigonella* L. *Cytologia* 48: 771-779.
- Amadi JE and Adeniyi DO (2009) Mycotoxin production by fungi isolated from stored grains. *Afr J Biotechnol* 8: 1219-1221.
- Angioi SA, Desiderio F, Rau D, Bitocchi E, Attene G and Papa R (2009) Development and use of chloroplast microsatellites in *Phaseolus* spp. and other legumes. *Plant Biol* 11:598-612.
- Arabi Z and Sardar S (2010) An investigation into the antifungal property of Fabaceae using bioinformatics tools. *Avicenna J Med Biotech* 2: 93-100.
- Arroyo-García R, Lefort F, María De Andrés T, Ibáñez J, Borrego J, Jouve N, et al.(2002) Chloroplast microsatellite polymorphisms in *Vitis* species. *Genome* 1149: 1142-1149.
- Avato P, Bucci R, Tava A, Vitali C, Rosato A, Bialy Z and Jurzysta M (2006) Antimicrobial activity of saponins from *Medicago* spp. : structure-activity relationship. *Phytother Res* 20:454-457.
- Bailly X, Olivieri I, Brunel B, Cleyet-Marel JC and Béna G (2007) Horizontal gene transfer and homologous recombination drive the evolution of the nitrogen-fixing symbionts of *Medicago* species. *J Bacteriol* 14: 5223-5236.
- Baldwin BG (1992) Phylogenetic utility of the internal transcribed spacers of nuclear ribosomal DNA in plants: An example from the Compositae. *Mol Phylogenet Evol* 1: 3–16.
- Baldwin BG, Sanderson MJ, Porter JM, Wojciechowski MF, Campbell CS and Donoghue MJ (1995) The ITS region of nuclear ribosomal DNA: a valuable source of evidence of angiosperm phylogeny. *Ann Mo Bot Gard* 82: 247-277.
- Bandelt HJ, Forster P and Röhl A (1999) Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol* 16:37-48.

- Baum BR (1968) A clarification of generic limits of *Trigonella* and *Medicago*. *Can J Bot* 46:741-749.
- Beckstrom-Sternberg SM and Duke JA (1994) The phytochemical database. <http://probe.nalusda.gov:8300/cgi-bin/query?dbgroup=phytochemdb> (ACEDB version 4.0 - data version July 1994).
- Bell LW, Ryan MH, Bennet RG, Collins MT and Clarked HJ (2012) Growth, yield and seed composition of native Australian legumes with potential as grain crops. *J Sci Food Agric* 92: 1354–1361.
- Bena G, Jubier MF, Olivieri I and Lejeune B (1998) Ribosomal external and internal transcribed spacers: combined use in the phylogenetic analysis of *Medicago* (Leguminosae). *J Mol Evol* 46: 299-306.
- Bena G (2001) Molecular phylogeny supports the morphological based taxonomic transfer of the “medicagoid” *Trigonella* species to the genus *Medicago* L. *Plant Syst Evol* 229: 217-236.
- Beneytout JL, Nappez C, Leboutet MJ and Malivvand G (1995) A plant steroid, diosgenin is a new meghakaryocytic differentiation inducer of HELA cell. *Biochem Biophys Res Comm* 207:398-404.
- Bertioli DJ, Seijo G, Fabio O, Freitas FO, Valls FM, Soraya CM, Bertioli L and Moretzsohn MC (2011) An overview of peanut and its wild relatives. *Plant Genet Res* 9: 134-149.
- Bhalodia NR and Shukla VJ (2011) Antibacterial and antifungal activities from leaf extracts of *Cassia fistula* L.: An ethnomedicinal plant. *J Adv Pharm Technol Res* 2: 104–109.
- Bhattarai N and Karki M. Medicinal and aromatic plants– Ethnobotany and conservation status. In: Burley J, Evans J and Youngquist J eds. *Encyclopedia of Forest Sciences*. Academic Press, London, UK, 2004, pp 523-532.
- Bhatti MA, Khan MTJ, Ahmed B and Jamshaid M (1996) Antimicrobial activity of *Trigonella foenum-graecum* seeds. *Fitoterapia* 67: 372-374.
- Bohannon MB, Hagemann JW and Earle FR (1974) Screening seeds of *Trigonella* and three related genera for diosgenin. *Phytochemistry* 13: 1513-1514.
- Borris RP(1996) Natural products research: perspectives from a major pharmaceutical company. *J Ethnopharmacol* 51: 29-38.
- Bremer K (1994) Branch support and tree stability. *Cladistics* 10: 295– 304.
- Brenac P and Sauvaire Y (1996) Accumulation of sterols and steroidal sapogenins in developing fenugreek pods: possible biosynthesis *in situ*. *Phytochemistry* 41:415-422.
- Brink S (1995). Looking beyond beta carotene. U.S. News and World Report, Washington DC, pp 92–93.
- Brockwell J, Evans MC, Bowman AM and McInnes A (2010) Distribution, frequency of occurrence and symbiotic properties of the Australian native legume *Trigonella suavissima* Lindl. and its associated root-nodule bacteria. *The Rangeland Journal* 32: 395-406.

- Buer CS, Gloria K, Muday GK and Djordjevic MA (2007) Flavonoids are differentially taken up and transported long distances in Arabidopsis. *Plant Physiol* 145: 478–490.
- Butters TD, Dwek RA, Platt FM. New therapeutics for the treatment of glycosphingolipid lysosomal storage diseases in Glycobiology and Medicine. In: Axford JS, ed. *Advances in Experimental Medicine and Biology*. New York Academic, 2003, pp 219-226.
- Bylka W, Matlawska I and Pilewski NA (2004) Natural flavonoids as antimicrobial agents. *J Am Nutraceut Assoc* 7: 24-31.
- Cai Z, Guisinger M, Kim HG, Ruck E, Blazier JC, McMurtry V *et al.* (2008) Extensive reorganization of the plastid genome of *Trifolium subterraneum* (Fabaceae) is associated with numerous repeated sequences and novel DNA insertions. *J Mol Evol* 67: 696–704.
- Cameron KM, Wurdack KJ and Jobson RW (2002) Molecular evidence for the common origin of snap-traps among carnivorous plants. *Am J Bot* 89: 1503–1509.
- Cannon SB, May GD and Jackson AS (2009) Three sequenced legume genomes and many crop species: rich opportunities for translational genomics. *Plant Physiol* 151: 970-977.
- Cannon SB, Sterck L, Rombauts S, Sato S, Cheung F, Gouzy J, Wang X, Mudge J, Vasdewani J, Schiex T, *et al.* (2006) Legume genome evolution viewed through the *Medicago truncatula* and *Lotus japonicus* genomes. *Proc Natl Acad Sci USA* 103: 14959–14964.
- Çeter T, Pinar M, Akan H, Ekici M and Aytaç Z (2012) Comparative seed morphology of *Trigonella* L. species (Leguminosae) in Turkey. *Afr J Agri Res* 7: 509-522.
- Chandra A (2011) Use of EST database markers from *M. truncatula* in the transferability to other forage legumes. *J Environ Biol* 32: 347-354.
- Chapagain B and Wiesman Z (2005) Variation in diaosgenin levels in seed kermels among different provenances of *Balanites aegyptiaca* Del (Zygophyllaceae) and its correlation with oil content. *Afr J Biotechnol* 4: 1209-1213.
- Chase MW, De AY, Buijn AV, Cox G, Reeves PJ, Rudall MAT, Johnson L and Eguiarte E (2000) Phylogenetics of Asphodelaceae Asparagales: an analysis of plastid *rbcL* and *trnL-F* DNA sequences. *Ann Bot* 86: 935-951.
- Chen PS, Shih YW, Huang HC and Cheng HW (2011) Diosgenin, a steroidal saponin, inhibits migration and invasion of human prostate cancer PC-3 Cells by reducing matrix metalloproteinases expression. *PLoS ONE* 6: e20164. doi:10.1371/journal.pone.0020164.
- Choudhary MI and Rehman A (2002) Recent Discoveries in Natural Products Chemistry, 7th EURASS. A conference on chemical Sciences, HEJ - Res. Inst. Chem., University of Karachi, Pakistan, pp: 25.
- Choudhary M and Yuvraj Gulia Y (2011) *Cassia Tora*: Its Chemistry, Medicinal Uses And Pharmacology. *Pharmacologyonline* 3: 78-96.
- Chu KT, Liu KH and Ng TB (2003) Cicerarin, a novel antifungal peptide from the green Chickpea. *Peptides* 24: 659-663.
- Compain P and Marti OR (2003) Design, synthesis and biological evaluation of iminosugar-based glycosyltransferase inhibitors. *Curr Top Med Chem* 3: 541-560.

- Cowan MM (1999) Plant Products as Antimicrobial Agents. *Clin MicroBiol Rev* 12: 564–582.
- Cuenca A, Escalante E and Pinero D (2003) Long-distance colonization, isolation by distance, and historical demography in a relictual Mexican pinyon pine (*Pinus nelsonii* Shaw) as revealed by paternally inherited genetic markers (cpSSRs). *Mol Ecol* 12: 2087–2097.
- Cunningham CW (1997a) Can three incongruence tests predict when data should be combined? *Mol Bio Evol* 14: 733–740.
- Cunningham CW (1997b) Is congruence between data partitions a reliable predictor of phylogenetic accuracy? Empirically testing an iterative procedure for choosing among phylogenetic methods. *Syst Biol* 46: 464–478.
- Dangi RS, Lagu MD, Choudhary LB, Ranjekar PK and Gupta VS (2004) Assessment of genetic diversity in *Trigonella foenum-graecum* and *Trigonella caerulea* using ISSR and RAPD markers. *BMC Plant Biology* 4:13.
- Dash BK, Sultana S and Sultana N (2011) Antibacterial activities of methanol and acetone extracts of Fenugreek (*Trigonella foenum*) and Coriander (*Coriandrum sativum*). *Life Sciences and Medicine Research* 1-8.
- De M, De AK and Banerjee AB (1999) Antimicrobial screening of some Indian spices. *Phytother Res* 13: 616-618.
- Dice LR (1945) Measures of the amount of ecologic association between species. *Ecology* 26: 297-302.
- Djordjevic MA, Redmond JW, Batley M and Rolfe BG (1987) Clovers secrete specific phenolic compounds which either stimulate or repress *nod* gene expression in *Rhizobium trifolii*. *EMBO J* 6: 1173-1179.
- Doebley J. Molecular systematics and crop evolution. In Soltis PS *et al.* eds. *Molecular Systematics of Plants*. Chapman & Hall, 1992, pp 202–222.
- Downie SR, Katz-Downie DS, Rogers EJ, Zujewski HL and Small E (1998) Multiple independent losses of the plastid *rpo* C1 intron in *Medicago* (Fabaceae) as inferred from phylogenetic analysis of nuclear ribosomal DNA internal transcribed spacer sequences. *Can J Bot* 76: 791 – 803.
- Doyle JJ (1992) Gene trees and species trees: molecular systematics as one-character taxonomy. *Syst Biol* 17: 144–163.
- Doyle JJ, Doyle JL, and Brown HD (1990) A chloroplast DNA phylogeny of the wild perennial relatives of soybean (*Glycine* subgenus *Glycine*): congruence with morphological and crossing groups. *Evolution* 44:371–389.
- Doyle JJ, Doyle JL, Ballenger JA, Dickson EE, Kajita T and Ohashi H (1997) A phylogeny of the chloroplast gene *rbcL* in the Leguminosae: taxonomic correlations and insights into the evolution of nodulation. *Am J Bot* 84: 541-554.
- Duangdeun M and Geoffrey C (1988) Traditional medicinal plants of Thailand XIII. Flavonoid derivatives from *Dracaena Loureiri* (AGAVACEAE). *J Sci Soc Thailand* 14: 3-21.
- Duke JA and Ayensu ES (1985) Medicinal Plants of China. Volume I and II, Reference publications. Inc.: United states of America. ISBN 0-917256-20-4.

- Dynowska M (1998) Hialohyphomycosis of the shank caused by *Fusarium solani*. *Med Mycol* 5: 241-245.
- Eliana-Bergter EB, Pinto MR, Rodrigijes ML. Structural and functional aspects of fungal glycosphingolipids. In: Atta-ur-Rahman, ed. *Studies in Natural Products Chemistry Vol 33, Part M Elsevier*, 2006, pp 1025-1055.
- Elmeer K and Almalki A (2011) DNA finger printing of *Prosopis cineraria* and *Prosopis juliflora* using ISSR and RAPD techniques. *Am J Plant Sci.* 2: 527-534.
- Ellison NW, Aaron A, JeVrey J, Steiner C, Williams WM and Taylor NL (2006) Molecular phylogenetics of the clover genus (*Trifolium*—Leguminosae). *Mol Phylogenet Evo* 39 :688–705.
- Elouafi I and Nachit MM (2004) A genetic linkage map of the Durum × *Triticum dicoccoides* L backcross population based on SSRs and AFLP markers, and QTL analysis for milling traits. *Theor Appl Genet* 108: 401–413.
- Facciola S (1990) *A Source Book of Edible Plants*. Kampong Publications.
- Farris JS (1989) The retention index and homoplasy excess. *Syst Zoo* 38: 406 – 407.
- Farris JS, Llersjo MKA, Kluge AG and Bult CJ (1994) Testing significance of incongruence. *Cladistics* 10: 315–319.
- Farris JS, Llersjo MKA, Kluge AG and Bult CJ (1995) Constructing a significance test for incongruence. *Syst Biol* 44:570–572.
- Farrukh A and Ahmad I (2003) Broad spectrum antibacterial and antifungal activity of certain traditionally used Indian medicinal plants. *World J Microbiol Biotechnol* 19: 653-657.
- Fazli FRY (1967) *Studies in steroid-yielding plants of the genus Trigonella*, PhD Thesis, University of Nottingham, England.
- Fazil FRY and Hardman R (1971) Isolation and characterization of steroids and other constituents from *Trigonella foenum-graecum*. *Phytochemistry* 10: 2497-2503.
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791.
- Fluit AC, Wienders LC, Verhoef J and Schmitz FJ (2001) Epidemiology and Susceptibility of 3,051 *Staphylococcus aureus* isolates from 25 University Hospitals participating in the European SENTRY study. *J Clin Microbiol* 39: 3727-3732.
- Flythe M and Kagan I (2010) Antimicrobial effect of red clover (*Trifolium pretense*) phenolic extract on the ruminal hyper ammonia producing bacterium, *Clostridium sticklandii*. *Curr Microbiol* 61: 125-131.
- Gao AG, Hakimi SM, Mittanck CA, Wu Y, Woerner BM, Stark DM, et al. (2000) Fungal pathogen protection in potato by expression of a plant defensin peptide. *Nat. Biotech* 18:1307-1310.
- Gatesy J and Baker R (2005) Hidden likelihood support in genomic data: can forty five wrongs make a right? *Syst Biol* 54: 483–492.

- Gazara M, Kamel W and Haider A (2001) Cladistic analysis of the genera: *Trifolium*, *Trigonella* and *Melilotus* (Fabaceae: Papilionaceae) in Egypt. *Egyptian Journal of Biology* 3: 161-170.
- Gepts P, Beavis WD, Brummer EC, Shoemaker RC, Stalker HT, Weeden NF and Young ND (2005) Legumes as a model plant family. In Genomics for food and feed. *Plant Phys* 137: 1228-1235.
- Ghislain M, Zhang D and Herrera MR (1999) Molecular biology laboratory protocols: plant genotyping training manual, 2<sup>nd</sup> edn. International Potato Center (CIP), Lima, Peru.
- Gibbons S (2004) Anti-staphylococcal plant natural products. *Nat Prod Rep* 21: 265-277.
- Gielly L and Taberlet P (1994) The use of chloroplast DNA to resolve phylogenies: non-coding versus *rbcl* sequences. *Mol Biol Evol* 11: 769-777.
- Graham PH and Vance CP (2003) Legumes: importance and constraints to greater use. *Plant Physiol* 131: 872-877.
- Greimel P, Spreitz J, Stütz AE and Wrodnigg TM (2003) Iminosugars and relatives as antiviral and potential anti-infective agents. *Curr Top Med Chem* 3: 513-518.
- Greuter W and Raus, Th. (ed.) 1987: Med-Checklist Notulae, 14. *Willdenowia* 16: 439-452.
- GRIN Taxonomy (2001) The Germplasm Resources Information Network. <http://www.ars-grin.gov/npgs/tax/taxgenform.html>
- Gunn CR and Kluve J (1976) Androecium and pistil characters for tribe Viciae (Fabaceae). *Taxon* 25:563-575.
- Guo X, Castillo-Ramirez S, Gonzalez V, Bustos P, Fernández-Vázquez JL, Santamaría R *et al.* (2007) Rapid evolutionary change of common bean (*Phaseolus vulgaris* L) plastome, and the genomic diversification of legume chloroplasts. *BMC Genomics* 8: 218. 228.
- Hall JB and Walker DH (1991) *Balanites aegyptiaca* Del. - A Monograph. School of Agricultural and Forest Sciences Publication, University of Wales, No. 3.
- Haouala R, Ouerghemml S, Tarchoune A and Boughanml N (2009) Improvement of *Trigonella maritima* Delile ex Poir. germination by polyploidization. *Pak J Bot* 41: 3001-3008.
- Hardman R and Fazil FRY (1972) Methods of screening the genus *Trigonella* for steroidal sapogenins. *Planta Medica* 21: 131-138.
- Hayes AJ and Markovic B (2002) Toxicity of Australian essential oils backhouse citriodora (lemon myrtle). Part-1. Antimicrobial activity and the in vitro cytotoxicity. *Food Chem Toxicol* 40: 535-543.
- Hector JN. Introduction to the Botany of Field Crops (Non cereals), Central News Agency Ltd., Johannesburg, 1936, pp 1127.
- Heyn CC. Trifolieae. In: Polhill RM and Raven PH, eds. Advances in Legume Systematics, Part 1. Royal Botanic Gardens, Kew, UK, 1981, pp 383-385.
- Hillis DM (1995) Approaches for assessing phylogenetic accuracy. *Syst Bio* 44: 3-16.

- Howie J, Ballard RA, De Koning C, Sandral G and Charman N (2001) *Trigonella balansae* – a new pasture legume for the alkaline soils of southern Australia? In: Proceedings of the 10<sup>th</sup> Australian Agronomy Conference, Hobart. Australian Society of Agronomy. Published online (<http://www.regional.org.au/au/asa/2001/3/d/howie.htm>).
- Hu J, Lavin MM, Wojciechowski MF and Sanderson MJ (2000) Phylogenetic systematics of the tribe Millettieae (Leguminosae) based on *matK* sequences and implications for evolutionary patterns in Papilionoideae. *Am J Bot* 87: 418-430.
- Huber-Morath. A *Trigonella* L. In: Davis PH, ed. Flora of Turkey and the East Aegean Islands. Edinburgh Univ. Press, Edinburgh, 1970, pp 452-482.
- Huelsenbeck JP, Larget B, Miller RE and Ronquist F (2002) Potential applications and pitfalls of Bayesian inference of phylogeny. *Syst Biol* 51, 673–688.
- Hungria M, Johnston AWB and Phillips DA (1992) Effects of flavonoids released naturally from Bean (*Phaseolus vulgaris*) on the *nod*-regulated gene transcription in *Rhizobium leguminosarum* *by. phaseoli*. *Mol Plant Microbe Interact* 5: 199-203.
- Hussain M, Hanif M, Ali S, Butcher R, Mirza T, Vanderveer D and Aziz-ur-Rehman (2010) *In vitro* biological studies and structural elucidation of fluoro-substituted phenyl acrylic acids. *Drug Dev Ind Pharm* 36: 1079 -1087.
- Hutchinson J. The Genera of Flowering Plants. Clarendon Press, Oxford. Vol. 1. 1964, pp 947 – 52.
- Ingham JL (1981) Phytoalexin induction and its chemosystematic significance. *Biochem Syst Ecol* 9: 275–81.
- Isely D (1998) Native and naturalized Leguminosae (Fabaceae) of the United States. Monte L. Bean Life Science Museum, Brigham Young University, Provo, Utah.
- Ishii T and McCouch SR (2000) Microsatellites and microsynteny in the chloroplast genomes of *Oryza* and eight other Gramineae species. *Theo Appl Gene* 100: 1257–1266.
- Jan WM, Linden DV, Warris A and Verweij EP (2011) *Aspergillus* species intrinsically resistant to antifungal agents *Med Mycol* 49:82–89.
- Janson RK, Wojciechowski MF, Sanniyasi E, Lee SB and Daniell H (2008) Complete plastid genome sequence of the chickpea (*Cicer arietinum*) and the phylogenetic distribution of *rps12* and *clpP* intron losses among legumes (Leguminosae). *Mol Phylogenet Evol* 48: 1204–17.
- Jaydev R, Jagan MRP, Malisetty V and Chinthalapally VR (2004) Diosgenin, a steroid saponin of *Trigonella foenum-graecum* (fenugreek), inhibits azoxymethane induced aberrant crypt foci formation in F344 rats and induces apoptosis in Ht-29 human Colon Cancer Cells. *Cancer Epidemiology Biomarkers and prevention* 13: 1392-1398.
- Jung HK, Eun-Soo P, Jae Hun S, Mal-Nam K, Woong-Sik M, Kyoo-Hyun C and Jin-San Y (2004) Antimicrobial Activity of *p*-Hydroxyphenyl acrylate derivatives. *J Agric Food Chem* 52: 7480–7483.
- Kakani RK, Singh SK, Pancholy A, Meena RS, Pathak R and Raturi A (2011) Assessment of genetic diversity in *Trigonella foenum-graecum* based on nuclear ribosomal DNA, Internal Transcribed Spacer and RAPD analysis. *Plant Mol Biol Rep* 29:315-323.



- Kan A, Özçelik B, Kartal M, Özdemir ZA and Özgen S (2010) *In Vitro* antimicrobial activities of *Cicer arietinum* L (Chickpea). *Tropical Journal of Pharmaceutical Research* 9: 475-481.
- Käss E and Wink M (1996) Molecular evolution of the Leguminosae: phylogeny of the three subfamilies based on *rbcL* sequences. *Biochem Sys Ecol* 24: 365-378.
- Käss E and Wink M (1997) Phylogenetic relationships in the Papilionoideae (Family Leguminosae) based on nucleotide sequences of cpDNA (*rbcL*) and nr DNA (ITS1 and 2). *Mol Phylo Evo* 8: 65-88.
- Kato T, Kaneko T, Sato S, Nakamura Y and Tabata S (2000) Complete structure of the chloroplast genome of a legume, *Lotus japonicas*. *DNA Res* 7: 323-30.
- Khan AV, Ahmed QU, Shukla I and Khan AA (2012) Antibacterial activity of leaves extracts of *Trifolium alexandrinum* Linn. against pathogenic bacteria causing tropical diseases. *Asian Pacific Journal of Tropical Biomedicine* 189-194.
- Khanna P and Jain SC (1973) Diosgenin, gitogenin and tigogenin from *Trigonella foenum-graecum* tissue cultures. *Lloydia* 36: 96-101.
- Kim KJ, Yu HH, Jeong SI, Cha JD, Kim SM and You YO (2004). Inhibitory effects of *Caesalpinia sappan* on growth and invasion of methicillin-resistant *Staphylococcus aureus*. *J Ethnopharmacol* 91:81-7.
- Kitiki A. Priority setting for underutilized and neglected plant species of Mediterranean region. In: Padulosi S, ed. Medicinal and aromatic plant species: opportunities for the Mediterranean region. ICARDA, Aleppo, Syria, 1998, pp 47-54.
- Kluge AG and Farris JS (1969) Quantitative phyletics and the evolution of anurans. *Sys Zoo* 8:1-32.
- Kores PJ, Molvray M, Weston PH, Hopper SD, Brown AP, Cameron KM and Chase W (2001) A phylogenetic analysis of *Diurideae* (Orchidaceae) based on plastid DNA sequence data. *Am J Bot* 88: 1903-1914.
- Kosslak RM, Bookland R, Barkei J, Paaren HE and Applebaum ER (1987) Induction of *Bradyrhizobium japonicum* common *nod* genes by isoflavones isolated from *Glycine max*. *Proc Natl Acad Sci USA* 84: 7428-7432.
- Kumar V, Srivastava N, Singh A, Vyas MK, Gupta S, Katudia K, et al. (2012) Genetic diversity and identification of variety-specific AFLP markers in fenugreek (*Trigonella foenum-graecum*). *Afr J Biotechnol* 11: 4323-4329.
- Küpper JM, Chioratto AF, Zucchi MI, Colombo CA, Carbonell SA, Mondego JM, et al. (2011) Genetic diversity in cultivated carioca common beans based on molecular marker analysis. *Genet Mol Biol* 34: 88-102.
- Joudi L and GH (2010) Exploration of medicinal species of Fabaceae, Lamiaceae and Asteraceae families in Ilkhji region, Eastern Azerbaijan Province (Northwestern Iran). *J Med Plants Res* 4: 1081-1084.
- Ladizinsky G (1979) Seed protein electrophoresis in the section *Foenum-graecum* of *Trigonella* L. (Fabaceae). *Pl Syst Evol* 133: 87-94.
- Lashin GMA (2006) Comparative morphology of pollen grains of some taxa of tribe Trifolieae (Fabaceae: Papilionoideae) from Egypt. *International Journal of Botany* 2: 270-277.

- Lavin M, Doyle JJ and Palmer JD (1990) Evolutionary significance of the loss of the chloroplast-DNA inverted repeat in the Leguminosae subfamily Papilionoideae. *Evolution* 44: 390–402.
- Lewis G, Schrire B, Mackind B, Lock M, eds. Legumes of the World. Royal Botanic Gardens, Kew, UK, 2005, pp 577.
- Li S, Du L, Yuen G and Harris SD (2006) Distinct ceramide synthases regulate polarized growth in the filamentous fungus *Aspergillus nidulans*. *Mol Biol Cell* 17: 1218–1227.
- Liang BB and Wang R (2004) The adverse reaction and advances in study on antifungal drugs. *Clin Med J* 2: 5-12.
- Lin S, Wu H, Jia H, Zhang P, Dixon R, May G, Gonzales R and Roe BA (2007) *Medicago truncatula* Variety Jema Long A-17 Chloroplast, Complete Sequence. Unpublished data GenBank number AC 093544.
- Liston A. Use of the polymerase chain reaction to survey for the loss of the inverted repeat in the legume chloroplast genome. In: Crisp MD and Doyle JJ, eds. Advances in legume systematic, part 7: Phylogeny Royal Botanic Gardens, Kew, 1995, pp 31–40.
- Liston A and Wheeler JA (1994) The phylogenetic position of the genus *Astragalus* (Fabaceae): evidence from the chloroplast genes *rpoC1* and *rpoC2*. *Biochem Syst Ecol* 22: 377–388.
- Liston A, Sterner J, Taylor NL, Ellison NW and Williams WM (2001) Phylogeny of New World *Trifolium*. Identification of a cryptic clade (abstract). 4<sup>th</sup> International Legume Conference.
- Liu FR and Miyamoto MM (1999) Phylogenetic assessment of molecular and morphological data for Eutherian mammals. *Syst Biol* 48: 54–64.
- Lock JM and Simpson K. Legumes of West Asia- a check-list. Royal Botanic Gardens, Kew, 1991, pp 263.
- Loi A, Nutt BJ, McRobb R and Ewing MA (2000). Potential new alternative annual pasture legumes for Australian Mediterranean farming systems. *Option Mediterranennes* 45: 51-54.
- Lust JB. The herb book. Bantam Books, New York, 1986, pp 1–55.
- Mabberley DJ. The plant book, 2nd ed. Cambridge University Press, Cambridge, UK, 1997.
- Mahesh B and Satish S (2008) Antimicrobial activity of some important medicinal plants against human pathogen. *World J Agri Sci* 4: 839-843.
- Malhotra SK. Fenugreek (*Trigonella foenum-graecum*L.). In: Sharma SK, ed. Breeding field crops. Academic Publisher, 2003, pp 802-835.
- Malhotra SK and Vashishtha BB (2008) Organic production of seed spices. National Research Centre for Seed Spices, Ajmer, India, pp 90.
- Mandal P, Sinha B and Mandal NC (2005) Antimicrobial activity of saponins from *Acacia auriculiformis*. *Fitoterapia* 76: 462-465.

- Marcilla A, Bargues MD, Ramsey JM, Magallon-Gastelum E, Salazar-Schettino P, Abad-Franch F, *et al.* (2001) The ITS-2 of the nuclear rDNA as a molecular marker for populations, species, and phylogenetic relationships in Triatominae (Hemiptera: Reduviidae), vectors of Chagas disease. *Mol Phy Evol* 18: 136-142.
- Marzougui N, Boubaya A, Elfalleh W, Guasmi F, Laaraiedh L, Ferchichi, Triki T and Beji M (2009) Assessment of genetic diversity in *Trigonella foenum graecum* Tunisian cultivars using ISSR markers. *J Food Agri Enviro* 7:101-105.
- Maureira-Butler IJ, Pfeil BE, Muangprom A, Osborn TC and Doyle JJ (2008) The reticulate history of *Medicago* (Fabaceae). *Syst Biol* 57: 466–482.
- Maxwell CA, Hartwig UA, Joseph CM and Phillips DA (1989) A chalcone and two related flavonoids released from alfalfa roots induce *nod* gene of *Rhizobium meliloti*. *Plant Physiol* 91: 842-847.
- Mayayo E, Pujol I and Guarro J (1999) Experimental pathogenicity of four opportunist *Fusarium* species in a murine model. *J Med Microbiol* 48: 363-366.
- McNeill J, Barrie FR and Burdet HM (2006) International code of botanical nomenclature (Vienna Code) adopted by the seventeenth International Botanical Congress, Vienna, Austria, July 2005 (electronic ed.), Vienna: International Association for Plant Taxonomy, <http://ibot.sav.sk/icbn/main.htm>, retrieved 2011-02-20, Article 19.7.
- Meepagala KM, Sturtz G and Wedge DE (2002) Antifungal constituents of the essential oil fraction of *Artemisia drancunculus* L. var. *Dracunculus*. *J Agri Food Chem* 50: 6989–6992.
- Mehrafarin A, Rezazadeh S, Badi N, Noormohammadi G, Zand E and Qaderi A (2011) A review on biology, cultivation and biotechnology of Fenugreek (*Trigonella foenum-graecum* L.) as a valuable medicinal plant and multipurpose. *J Med Plant* 10: 6-24.
- Mengoni A, Ruggini C, Vendramin VV and Bazzicalupo M (2000) Variation in tetraploid alfa-alfa. *Plant Breed* 119: 509 -515.
- Merklin A, Christen P and Kapetanidis I (1997) Production of diosgenin by hairy root cultures of *Trigonella foenum.graecum* L. *Plant Cell Rep* 16:632-636.
- Moalic S, Liagre B and Corbiere C (2001) A plant steroid, diosgenin induces apoptosis, cell cycle arrest and cox activity in osteosarcoma cells. *FEBS Lett.* 506: 225-30.
- Mohana DC and Raveesha KA (2006) Antibacterial activity of *Caesalpinia coriaria* (Jacq.) Willd. against plant pathogenic *Xanthomonas* pathogens: an ecofriendly approach. *J Agri Technol* 2: 317-327.
- Moore WS (1995) Inferring phylogenies from mtDNA variation: mitochondrial-gene trees versus nuclear-gene trees. *Evolution* 49: 718–726.
- Morrissey JP and Osbourn AE (1999) Fungal resistance to plant antibiotics as a mechanism of pathogenesis. *Microbiol Mol Biol Rev* 63: 708-724.
- Morrissey JP, Wubben JP and Osbourn AE (2000) *Stagonospora avenae* secretes multiple enzymes that hydrolyze oat leaf saponins. *Mol Plant Microbe Interact* 13: 1041–1052.

- Muehlbauer FJ. Use of wild species as a source of resistance in cool-season food legume crops. In: Singh KB, Saxena MC, eds. Breeding for stress tolerance in cool-season food Legume; Aleppo: ICARDA , 1993, pp 359-372.
- Mulabagal V and Tsay HS (2004) Plant Cell Cultures - An alternative and efficient source for the production of biologically important secondary metabolites. *Int J Appl Sci Eng* 1: 29-48.
- Mulabagal V, Chen-yee-Lee, Lo SF, Nalawade SM, Lin CY and Tsay H (2003) Studies on production of some important secondary metabolites from medicinal plants by plant tissue culture. *Bot Bull Acad Sin* 45 1-22.
- Nair RM, Dundas IS, Wallwork M, Verlin DC, Waterhouse L and Dowling K (2004) Breeding system in a population of *Trigonella balansae* (Leguminosae). *Annu Bot* 94: 883–888.
- Niknam V, Sharifzadeh B, Ebrahimzadeh H and Zarre S (2004) Comparative study of proteins in seeds of some species of *Trigonella* from Iran. *Iranian Int J Sci* 5:1-11.
- Nixon KC and Carpenter JM (1996) On simultaneous analysis. *Cladistics* 12: 221–241.
- Novák K, Chovanec P, Škrdleta V, Kropáčová M, Lisá L and Němcová M (2002) Effect of exogenous flavonoids on nodulation of pea (*Pisum sativum* L.). *J Exp Bot* 53: 1735-1745.
- Oguyemi AO. Phytochemical Methods. In: Sofowora A, ed. Proceedings of a Conference on African Medicinal Plants. Ife-Ife: Univ Ife; Harbone JB. London: Chapman and Hill, 1979, pp 20-22.
- Olli S and Kirti PB (2006) Cloning, characterization and antifungal activity of defensin Tfgd1 from *Trigonella foenum-graecum* L. *J Biochem and Mol Bio* 39: 278-283.
- Oncina R, Botia JM, Rio JA and Ortuno A (2000) Bioproduction of diosgenin in callus cultures of *T. foenum-graecum* L. *Food chemistry* 70: 489-492.
- Oncina R, Botia JM, Rio JA and Ortuno A (2002) Effect of ethylene on diosgenin accumulations in the callus cultures of *T. foenum-graecum* L. *Food chemistry* 76: 475-479.
- Ortuno A, Oncina R, Botia JM and Rio JA (1998) Distribution and changes of diosgenin during development of *T. foenum-graecum* plants. Modulation by benzylaminopurine. *Food chemistry* 63: 51-54.
- Ortuno A, Oncina R, Botia JM and Rio JA (1999) Regulation of the diosgenin expression in *T. foenum-graecum* plants by different plant growth regulators. *Food chemistry* 65: 227-232.
- Osborn TC, Brouwer D and McCoy TJ. Molecular marker analysis of alfalfa. In McKersie BD and Brown DCW, eds. Biotechnology and the improvement of forage legumes, 1997, pp 91-109.
- Osbourn A (1996) Saponins and plant defence—a soap story. *Trends Plant Sci* 1: 1-6.
- Pang X, Cong Y, Yu H, Kang L, Feng B, Han B *et al.* (2012) Spirostanol saponins derivatives from the seeds of *Trigonella foenum-graecum* by  $\beta$ -glucosidase hydrolysis and their inhibitory effects on rat platelet aggregation. *Planta Med* 78: 276–285.
- Parekh J and Chanda S (2006) *In vitro* antimicrobial activity and phytochemical analysis of some Indian medicinal Plants. *Turkish Journal of Biology* 31: 53-58.

- Parthasarathy VA, Kandiannan K and Srinivasan. Introduction. In Parthasarathy VA, Kandiannan K and Srinivasan V. eds. Organic spice, New India publishing agency, 2008, pp 1-38
- Pattnaik S, Padhan DK and Jana GK (2010) Evaluation of cinnamon oil, peppermint oil, cardamom oil and orange oil as antimicrobial agents. *J Pharm Res* 3: 414–416.
- Pepljnak S, Kalodeera Z and Zovko M (2005) Antimicrobial activity of flavonoids from *Pelargonium redula* (Cav.) L Herit *Acta Pharm* 55: 431-435.
- Perez C, Pauli M and Bazerqu P (1990) An antibiotic assay by the well agar method. *Acta Biol Med Exp* 15:113-115.
- Peters NK, Frost JW and Long SR (1986) A plant flavone, luteolin, induces expression of *Rhizobium meliloti* nodulation genes. *Science* 233: 977-980.
- Petropoulos GA (1973) Agronomic, genetic and chemical studies of *Trigonella foenum-graecum* L. PhD. Diss. Bath University, England. 145.
- Petropoulos GA. Fenugreek, The genus *Trigonella*. Taylor and Francis, London and New York, 2002, pp 1 - 255.
- Polhill RM. Classification of the Leguminosae. In: Bisby FA, Buckingham J and Harborne JB, eds. Phytochemical dictionary of the Leguminosae, xxxv–xlvi. Chapman and Hall, New York, 1994, USA.
- Posada D (2008) jModelTest: Phylogenetic Model Averaging. *Mol Biol and Evol* 25: 1253-1256.
- Posada D and Crandall KA (1998) Modeltest, testing the model of DNA substitution. *Bioinformatics* 14: 817-818.
- Powell W, Morgante M, Andre C, McNicol JW, Marchray GC, Doyle JJ, *et al.*(1995) Hypervariable microsatellites provide a general source of polymorphic DNA markers for the chloroplast genome. *Curr Bio* 5: 1023-1029.
- Powell W, Morgante M, Doyle J, McNicol JW, Tingey SV and Rafalski AJ (1996) Genepool variation in the genus *Glycine* subgenus *soja* revealed by polymorphic nuclear and chloroplast microsatellites. *Genetics* 144: 792–803.
- Prevost A and Wilkinson MJ (1999) A new system of comparing PCR primers applied to ISSR fingerprinting of potato cultivars. *Theor Appl Genet* 98: 107-112.
- Provan J, Corbett G, McNicol JW and Powell W (1997) Chloroplast variability in wild and cultivated rice (*Oryza* spp.) revealed by polymorphic chloroplast simple sequence repeats. *Genome* 40: 104–110.
- Provan J, Russell JR, Booth A and Powell W (1999) Polymorphic chloroplast simple-sequence repeat primers for systematic and population studies in the genus *Hordeum*. *Mol Ecol* 8: 505–511.
- Provan J, Powell W and Hollingsworth PM (2001) Chloroplast microsatellites: new tools for studies in plant ecology and evolution. *Trends in Eco and Evo* 16: 142–147.
- Puri HS, Jefferies TM and Harh R (1976) Diosgenin and yamogenin levels in some Indian plant samples. *Planta, Med.* 30: 118-121.

- Rajagopalan MS (1998) Fenugreek, what this herb can offer? *Naturally*. 1: 1-4.
- Rajpal V. Standardization of Botanicals (testing and extraction methods of medicinal herbs); Vol.1, Eastern Publishers, New Delhi, India; 2002, pp 39-46.
- Rechinger KH. *Trigonella*. In: Rechinger KH, ed. Flora Iranica no. 157: 200-253. -Graz: Akadem, 1984
- Recourt K, Schripsema J, Kijne JW, Brussel A and Lugtenberg BJ (1991) Inoculation of *Vicia sativa* subsp. *nigra* roots with *Rhizobium leguminosarum* bv. *viciae* results in release of *nod* gene activating flavanones and chalcones. *Plant Mol Biol* 16: 841-852.
- Rezaeian S (2011) Assessment of diosgenin production by *Trigonella foenum-graecum* L. *in vitro* condition. *Am J Plant Physiol* 6: 261-268.
- Rochfort S and Panozzo J (2007) Phytochemicals for health, the role of pulses. *J Agri Food Chem*. 55: 7981-7994.
- Rogers SO and Bendich AJ (1988) Extraction of DNA from milligram amounts of fresh, herbarium and mummified plant tissues. *Plant Mol Biol* 5: 69-76.
- Rohlf FJ (2000) *NTSYS-pc: Numerical Taxonomy System*. Ver. 2.1. Exeter Software, Setauket, NY, USA. pp. 29-34.
- Rokas A, Williams BL, King N and Carroll SB (2003) Genome-scale approaches to resolving incongruence in molecular phylogenies. *Nature* 425: 798-804.
- Roman ID, Thewles A and Coleman R (1995) Fractionation of livers following diosgenin treatment to elevate binary cholesterol. *Biochemistry Biophysics Acta* 1255: 77-81.
- Rongwen J, Akkaya MS, Bhagwat AA, Lavi U and Cregan PB (1995) The use of microsatellite DNA markers for soybean genotype identification. *Theor Appl Genet* 90:43-48.
- Ronquist F and Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572-1574.
- Rossen L, Davis EO and Johnston AWB (1987) Plant-induced expression of *Rhizobium* genes involved in host specificity and early stages of nodulation. *Trends Biochem Sci* 12: 430-433.
- Rouk HF and Mangesha H. Fenugreek (*Trigonella foenum-graecum* L. Its relationship, geography and economic importance, Exper Stat Bull No. 20, Imper. Ethiopian College of Agric. & Mech. Arts.
- Saleem M, Nazir M, Ali MS, Hussain H, Lee YS, Riaz N and Jabbar A (2010) Antimicrobial natural products: an update on future antibiotic drug candidates. *Nat Prod Rep* 27: 238-242.
- Saleh NAM, Boulos L, El-Negoumy SI and Abdalla MF (1982) A comparative study of the flavonoids of *Medicago radiata* with other *Medicago* and related *Trigonella* species. *Biochem Syst Ecol* 10: 33-6.
- Sambrook J and Russell DW. *Molecular Cloning: A Laboratory Manual*, Volume 3, Cold Spring Harbor Laboratory Press, 2001.

- Sanderson MJ and Wojciechowski MF (1996) Diversification rates in a temperate legume clade: Are there “so many species” of *Astragalus* (Fabaceae)? *Am J Bot* 83: 1488–1502.
- Sandor PK, Sandor MK and Andras K (2004) Comparative test of fenugreek/ *Trigonella-foenum-graecum* varieties. *Central European Agriculture*. 5: 259-262.
- Saski C, Lee SB, Daniell H, Wood TC, Tomkins J, Kim HG and Jansen RK (2005) Complete chloroplast genome sequence of *Glycine max* and comparative analysis with other legume genomes. *Plant Mol Biol* 59: 309–22.
- Satish S, Raveesha KA and Janardhana GR (1999) Antibacterial activity of plant extracts on phytopathogenic *Xanthomonas campestris* pathovars. *Lett Appl Microbiol* 28: 145-147.
- Sato S, Nakamura Y, Kaneko T, Asamizu E, Kato T, Nakao M, et al. (2008) Genome structure of the legume, *Lotus japonicus*. *DNA Res* 15: 227–239
- Savikin-Fodulovic K, Grubisic D, Culafic L, Menkovic N and Ristic M (1998) Diosgenin and phytosterols content in five callus lines of *Dioscorea balcanica*. *Plant Sci* 135:63-67.
- Schmidt C (2004) Analysis of 7'4'-Dihydroxyflavone by Positive APCI LC/MS/MS.CC-MS. *Varian, Inc* 12.
- Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, et al. (2010) Genome sequence of the palaeopolyploid soybean. *Nature* 14: 178-183.
- Schramm VL and Tyler PC (2003) Immune- sugar- based nucleosides. *Curr Top Med Chem* 3: 525-532.
- Schultz OF (1901) Monographie der Gattung *Melilotus*. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 29: 660 – 735.
- Seelanan T, Schnabel A and Wendel JF (1997) Congruence and consensus in the cotton tribe (Malvaceae). *Syst Biol* 22: 259–290.
- Sharma RD (1990) Effect of Fenugreek on blood glucose and serum lipids in type I diabetes. *European J Clinical Nutrition* 44:301–306.
- Shiva KN and Malhotra SK. Fenugreek. In: Parthasarathy VA, Kandiannan K and Srinivasan V, eds. Organic spice. New India publishing agency, 2008, pp 547-562.
- Simin K, Ali Z and Khaliq-Uz-Zaman SM (2002) Structure and biological activity of a new rotenoid from *Pongamia pinnata*. *Nat Prod Res* 16:351–357.
- Singh A and Saini RG (1971) Cytological studies on triploid *Trigonella corniculata* L. *Genetica* 42: 469-476.
- Singh A (1973) Studies on the interspecific hybrids of *Trigonella corniculata* L., *T. hamosa* L. and *T. cretica* L. *Genetica* 44: 264-269.
- Singh KN and Kaushal R (2007) Comprehensive notes on commercial utilization, characteristics and status of steroid yielding plants in India. *Ethnobotanical Leaflets* 11: 45-51.
- Singh NK, Gupta DK, Jayaswal PK, Mahato AK, Dutta S, Singh S et al. (2012) The first draft of the pigeonpea genome sequence *J Plant Biochem Biotechnol* 21: 98-112

- Sinskaya E. Flora of the Cultivated Plants of the U.S.S.R. XIII. In Perennial Leguminous Plants. Part I: Medicago, Sweet clover, Fenugreek. Israel Programme for Scientific Translations, Jerusalem, 1961, pp 514–70.
- Sirjaev G (1935) Die Entwicklungsgeschichte der Gattung *Trigonella* (Suppl. Ad Generis *Trigonella* revisio critica) *Bull. Assoc. Russe Rech Sci Prague* 2: 135-162.
- Skaltsa H. Chemical composition. In: Petropoulos GA, ed. Fenugreek, The genus *Trigonella*. Taylor and Francis, London and New York, 2002, pp 132-160.
- Small E (1987a) A taxonomic study of the “medicagoid” *Trigonella* (Leguminosae).
- Small E (1987b) Polythetic generic separation in the tribe Trifolieae subtribe Trigonellinae (Leguminosae). *Can J Bot* 67: 1480 – 1492.
- Small E. Generic changes in *Trifolieae* subtribe *Trigonellinae*. In: Stirton CH (ed) Advances in legume systematics, part 3, vol 3. Royal botanic gardens, Kew, 1987-c, pp169.
- Small E (1988) Pollen-ovule patterns in tribe Trifolieae (Leguminosae). *Plant Syst Evol* 160:195–205.
- Small E (2010) Alfalfa and relatives: Evolution and classification of *Medicago*. NRC Research Press, Ottawa, Ontario, Canada. Small E and Jomphe M (1988) A synopsis of the genus *Medicago* (Leguminosae). *Can J Bot* 67: 3260–3294.
- Small E and Brookes BS (1990a) A numerical taxonomic analysis of the *Medicago littoralis* - *M. truncatula* complex. *Can J Bot* 68: 1667 – 1674.
- Small E and Brookes BS (1990b) A taxonomic simplification of *Medicago italica*. *Can J Bot* 68: 2103 – 2111.
- Small E and Jomphe M (1989a) A note on *Trigonella* (*Melilotus*) *bicolor*. *Can J Bot* 67: 1604 – 1606.
- Small E and Jomphe M (1989b) A synopsis of the genus *Medicago* (Leguminosae). *Can J Bot* 67: 3260 – 3294.
- Small E, Crompton CW and Brookes BS (1981) The taxonomic value of the floral characters in the tribe Trigonellinae (Leguminosae), with special reference to *Medicago*. *Can J Bot* 59:1578-1598.
- Smýka IP, Hýbl M, Corander J, Jarkovský J, Flavell AJ and Griga M (2008) Genetic diversity and population structure of pea (*Pisum sativum* L.) varieties derived from combined retrotransposon, microsatellite and morphological marker analysis. *Theor Appl Genet* 117:413-424.
- Sorenson MD (1999) TreeRot, version 2. Boston University, Boston, Massachusetts.
- Sorkheh K, Shiran B, Gradziel TM, Epperson BK, Martýnez-Gomez P and Asadi E (2007) Amplified fragment length polymorphism as a tool for molecular characterization of almond germplasm: genetic diversity among cultivated genotypes and related wild species of almond, and its relationships with agronomic traits. *Euphytica* 156: 327-344.
- Spataro G, Tiranti B, Arcaleni P, Bellucci E, Attene G, Papa R, Spagnoletti PZ and Negri V (2011) Genetic diversity and structure of a worldwide collection of *Phaseolus coccineus* L. *Theor Appl Genet* 122:1281-1291.



- Steele KP and Wojciechowski MF (2003) Phylogenetic analyses of tribes Trifolieae and Viciae, based on sequences of the plastid gene *matK* (Papilionoideae: Leguminosae). In: Klitgaard BB and Bruneau A, eds. *Advances in Legume Systematics*, part 10, higher level systematics, Royal Botanic Garden, Kew, UK, 2003, pp 355–370.
- Steele KP, Ickert-Bond SM, Zarre S and Wojciechowski MF (2010) Phylogeny and character evolution in *Medicago* (Leguminosae): evidence from analyses of plastid *trnk/matk* and nuclear *GA3ox1* sequences. *Am J Bot* 97:1142–1155.
- Stütz AE. Iminosugars as Glycosidase Inhibitors: Nojirimycin and Beyond. New York: Wiley-VCH, 1999.
- Sucher NJ and Charles MC (2008) Genome based approaches for authentication of medicinal plants. *Planta Med* 74: 603-623.
- Sukhotu T, Kamijima O and Hosaka K (2006) Chloroplast DNA variation in the most primitive cultivated diploid potato species *Solanum stenotomum* Juz. et Buk. and its putative wild ancestral species using high-resolution markers. *Genet Resour Crop Evol* 53: 53–63.
- Sumitra C (2006) Screening of aqueous and alcoholic extracts of some Indian medicinal plants for antibacterial activity. *Indian J Pharm Sci* 68: 835-838.
- Sundaram S and Purwar S (2011) Assessment of genetic diversity among fenugreek (*Trigonella foenum-graecum* L.), using RAPD molecular markers. *J Med Plant Res* 5:1543-1548.
- Swaffer DS, Ang CY, Desai PB and Rosenthal GA (1994) Inhibition of the growth of human pancreatic cancer cells by the arginine antimetabolite L-canavanine. *Cancer Res* 54:6045–6048.
- Swaffer DS, Ang CY, Desai PB, Rosenthal GA, Thomas DA, Crooks PA and John WJ (1995) Combination therapy with 5-fluorouracil and L-canavanine: In vitro and in vivo studies. *Anti-cancer Drugs* 6:586–593.
- Swofford DL (2001) PAUP\*: phylogenetic analysis using parsimony (\*and other methods), version 4. Sinauer, Sunderland, Massachusetts, USA.
- Taberlet P, Gielly L, Pautou G and Bouvet J (1991) Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant Mol Biol* 17: 1105–1109.
- Tan A. Biodiversity conservation *ex situ*, *in situ* conservation: A case in Turkey. In: Watanabe K and Komamine A, eds. *Challenge of Plant and Agricultural Sciences to the crisis of biosphere on the Earth in the 21st Century*. Eurekah, Texas, 2000.
- Tangphatsornruang S, Sangsrakru D, Chanprasert J, Uthaipaisanwong P, Yoocha T, Jomchai N and Tragoonrung S (2010) The chloroplast genome sequence of mungbean (*Vigna radiata*) determined by high-throughput pyrosequencing: Structural organization and phylogenetic relationships. *DNA Res* 17: 11–22.
- Tantasawat P, Trongchuen J, Prajongjai T and Thongpae T (2010) Variety identification and genetic relationships of mungbean and blackgram in Thailand based on morphological characters and ISSR analysis. *Afr J Biotechnol* 9: 4452-4464.
- Tar'an B, Zhang C, Warketin T, Tullu A and Vandenberg A (2005) Genetic diversity among varieties and wild species accessions of pea (*Pisum sativum* L.) based on molecular markers and morphological and physiological characters. *Genome* 48: 257-272.

- Taylor WG, Zulynaik HJ, Richards KW, Acharya SN, Bittman S and Elder JL (2002) Variation in diosgenin levels among 10 accessions of fenugreek seeds produced in western Canada. *J Agri Food Chem* 50: 5994-5997.
- Thomas JE, Basu SK and Acharya SN (2007) Identification of *Trigonella* accessions which lack antimicrobial activity and are suitable for forage development. *Can J Plant Sci* 86: 727-732.
- Thomas MR, Matsumoto S, Cain P and Scott NS (1993) Repetitive DNA of grapevine: classes present and sequences suitable for cultivar identification. *Theor Appl Genet* 86:173-180.
- Thomma B, Cammue B and Thevissen K (2003) Mode of action of plant defensins suggests therapeutic potential. *Curr Drug Targets* 3: 1-8.
- Thompson JD, Higgins DG and Gibson TJ (1994) CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22: 4673-4680.
- Tim TP and Lamb AJ (2005) Antimicrobial activity of flavonoids. *IJOAA* 26: 343-356.
- Trivedi P (2007) A validated quantitative thin layer chromatography method for the estimation of diosgenin in various plant samples, extract and market formulations. *Journal of AOAC International* 90:358-363.
- Tutin TG and Heywood VH. Flora Europaea. Vol. I & II. Cambridge University Press, Cambridge. 1964, pp 452-482.
- Tyler VE, Brady LR and Robbers JE (1976) Glycosides. In: Pharmacognosy. Lea and Febiger, Philadelphia, PA, 7<sup>th</sup> Edition, 1976, pp 76-103.
- Varshney RK, Thudi M, Aggarwal RK and Börner A. Genic Molecular Markers in plant development and application. In: Varshneya R and Tuberosa R. eds. Genomics-Assisted Crop Improvement, Volume 1: Genomics Approaches and Platforms. Publisher Springer, 2007, pp13-31.
- Varshney RK and Thudi M (2010) Legume genomics and breeding. *Plant Breed Rev* 33: 257-304.
- Varshney RK, Chen W, Li Y, Bharati AK, Saxena RK, Schlueter IA, Donoghue MTA *et al.* (2012) Draft genome sequence of pigeonpea (*Cajanus cajan*), an orphan legume crop of resource – poor farmers. *Nature Biotechnology* 30:83-89.
- Vasil'chenko IT (1953). Bericht über die Arten der Gattung, *Trigonella* Trudy Bot. Inst. Akad. Nauk. *Flora of Russian*. 1:10.
- Wagh P, Rai M, Deshmukh SK and Durate MCT (2007) Bio-activity of oils of *Trigonella foenum-graecum* and *Pongamia pinnata*. *Afr J Biotechnol* 6: 1592-1596.
- Watson LE, Sayed-Ahmed H and Badr A (2000) Molecular phylogeny of Old World *Trifolium* (Fabaceae). *Pl Syst Evol* 224: 153-171.
- Weir BS (1996) Genetic data analysis II. *Sinauer Publisher, Sunderland, MA*.

- Weising K and Gardner RC (1999) A set of conserved PCR primers for the analysis of simple sequence repeat polymorphisms in chloroplast genomes of dicotyledonous angiosperms. *Genome* 42:9-19.
- White T J, Bruns T, Lee S and Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, and White TJ, eds. "PCR Protocols: A Guide to Methods and Applications", Academic Press, New York, 1990, pp 315–322.
- Wojciechowski MF. Reconstructing the phylogeny of legumes (Leguminosae): an early 21<sup>st</sup> century perspective. In: B.B; Klitgaard and Bruneau A, ed. Advances in legume systematic, part 10, Higher level systematic, Royal Botanic garden, Kew, 2003, pp 5-35.
- Wojciechowski MF, Lavin M and Sanderson MJ (2004) A phylogeny of legumes (Leguminosae) based on analysis of the plastid *matK* gene resolves many well-supported subclades within the family. *Am J Bot* 91: 1846–1862.
- Wojciechowski MF, Sanderson MJ, Steele KP and Liston A Molecular phylogeny of the "Temperate Herbaceous Tribes" of Papilionoid legumes: a supertree approach. In: Herendeen PS and Bruneau A, ed. Advances in Legume Systematics, *Royal Botanic Gardens, Kew*, 2000, pp 277–298.
- Xu DH, Abe J, Gai JY and Shimamoto Y (2002) Diversity of chloroplast DNA SSRs in wild and cultivated soybeans: evidence for multiple origins of cultivated soybean. *Theo Appl Genet* 105: 645–653.
- Yadav RN and Jain S (2005) A new bioactive flavone glycoside from the seeds of *Melilotus indica* All. *J Asian Nat Prod Res* 7: 595-599.
- Yap IV and Nelson RJ (1996) WinBoot: a program for performing bootstrap analysis of binary data to determine the confidence limits of UPGMA-based dendrograms. IRRRI Disc. Ser. No. 14. Int. Rice Res. Inst., Manila, Philippines.
- Ye XY, Wang HX and Ng TB (2000) Sativin, a novel antifungal miraculin-like protein isolated from legumes of the sugar snap *Pisum sativum* var. *macrocarpon*. *Life Sciences* 67: 775-781.
- Young ND, Debelle F, Oldroyd GE, Geurts R, Cannon SB, Udvardi MK, *et al.* (2011) The *Medicago* genome provides insight into the evolution of rhizobial symbioses. *Nature* 480: 520-524.
- Zaat AJS, Carel A, Wuffelman AC, Mulders HMI, Brussela AN and Lugtenberg JJB (1988) Root exudates of various host plants of *Rhizobium leguminosarum* contain different sets of inducers of *Rhizobium* nodulation genes. *Plant Physiol* 86: 1298-1303.
- Zuanazzi JAS, Clergeot PH, Quirion JC, Husson HP, Kondorosi A and Ratet P (1998) Production of *Sinorhizobium meliloti nod* gene activator and repressor flavonoids from *Medicago sativa* roots. *Mol Plant Microbe Interact* 11: 784–794.
- Zietkiewicz E, Rafalski A and Labuda D (1994) Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* 20:176–183.