

## List of Figures

### **Chapter 1:**

Fig. 1.1: *Trigonella*- type species *T. foenum-graecum*

Fig. 1.2: Herbal formulations containing *T. foenum-graecum*

Fig. 1.3: Fenugreek growing states in India

Fig. 1.4: Year wise production of fenugreek in India

Fig.1.5: (a) *T. caerulea*: blue fenugreek (b) Swiss cheese flavored with blue fenugreek

Fig. 1.6: *T. corniculata* - "Kasuri methi"

Fig 1.7: *T. balansae* - Upright growth habit and the proliferation of pods at the top of the canopy make harvesting of seeds easier.

Fig 1.8: *T. suavissima* - potential as a grain crop in arid and nutrient poor environments

Fig.1.9: IRLC - The Inverted Repeat-lacking clade (Wojciechowski *et al.* 2000)

Fig. 1.10: The position of *Trifolium*, *Medicago*, *Trigonella* and *Melilotus* among the genera of the "vicioid clade" (Ellison *et al.* 2006)

### **Chapter 2:**

Fig: 2.1: Variations in plant habit and leaf size of (a) *T. caerulea* EC 583569 (Spain) and ( b) *T. caerulea* EC 583568 (Turkey)

Fig: 2.2: Variations in seed size in (a) *T. caerulea* EC 583567 (Australia), (b) *T. caerulea* EC 583568 (Turkey) and (c) *T. caerulea* EC 583569 (Spain).

Fig. 2.3: Variation in seed size in (a) *T. filipes* EC 583580 (Syria), (b) *T. filipes* EC 583581 (Iran) and (c) *T. filipes* EC 583584 (Turkey).

Fig. 2.4: Variation in seed size in (a) *T. kotschyi* EC 583598 (Turkey) and (b) *T. kotschyi* EC 58397 (Jordan).

Fig. 2.5: Variations between *T. coelesyriaca* accessions, EC 583557 and EC 583562. a-b Comparison of plant height; c-d leaf margin and length; e-f pod shape and length; g-h seed morphology.

Fig. 2.6: Variation between *T. schlumbergeri* accessions EC 583609, EC 583611 and EC 583612. a-c Comparison of p leaf margin and length; d-f pod shape and length; g-i seed morphology.

### Chapter 3:

Fig 3.1: (a) Amplification profile obtained with ISSR primer UBC807 showing a species specific band in *T. foenum-graecum* accessions.

Fig 3.1: (b) Amplification profile obtained with ISSR primer UBC880 showing species specific bands in *T. cretica* accessions

Fig 3.2: (a) Amplification profile obtained with primer ccmp 10 showing intraspecific variation in accessions of *T. caerulea*

Fig 3.2: (b) Amplification profile obtained with primer ccmp 4 showing intraspecific variation in accessions of *T. cylindracea*, *T. filipes*, *T. kotschyi* and *T. mesopotamica*

Fig. 3.3: UPGMA based dendrogram showing relationships between *Trigonella* species revealed by ISSRs.

Fig. 3.4: UPGMA based dendrogram showing relationships between *Trigonella* species revealed by Cp SSRs.

Fig.3.5: Haplotype network for 22 *Trigonella* species.

### Chapter 4:

Fig. 4.1: Organization of the nr DNA region. Arrows indicate the approximate position of the primer site

Fig. 4.2: Position and direction of the universal primers used to amplify non-coding regions of chloroplast DNA

Fig. 4.3: Maximum parsimony analysis of combined ITS + *trnL-F* sequences.

Fig. 4.4: Intraspecific nucleotide sequence divergence observed in ITS region

Fig. 4.5: Intraspecific nucleotide sequence divergence observed in *trnL-F* region

Fig.4.6: Phylogenetic relationship in *Trigonella* based on the maximum parsimony analysis of ITS-I sequences

Fig.4.7: Phylogenetic relationship in *Trigonella* based on the maximum parsimony analysis of ITS-II sequences

Fig.4.8: Phylogenetic relationship in *Trigonella* based on the maximum parsimony analysis of ITS sequences.

Fig.4.9: Phylogenetic relationship in *Trigonella* based on the maximum parsimony analysis of *trnL-F* sequences.

Fig.4.10: Phylogenetic relationship in *Trigonella* based on the maximum parsimony analysis of ITS+ *trnL* intron sequences.

Fig.4.11: Phylogenetic relationship in *Trigonella* based on the maximum parsimony analysis of ITS+ *trnL-F* sequences.

## **Chapter 5:**

Fig. 5.1: Preliminary phytochemical analysis

Fig.5.2: Antibacterial activity of the crude methanol extract of some *Trigonella* species against (a) *S. aureus* and (b) *E. coli*

Fig.5.3: Antibacterial activity of the sapogenin extract against (a) *B. subtilis* and (b) *S. aureus*

Fig.5.4: Antibacterial activity of the polar and non-polar extract prepared from dried sapogenin extract.

Fig.5.5: Antibacterial activities of fractions collected from (a) first and (b) second silica gel column chromatography against *B. subtilis*

Fig.5.6: Total chromatogram of fraction 6 obtained at low cone voltage

Fig.5.7: Elemental composition of  $m/z=255.0653$  determined by MassLynx v4.1 software

Fig. 5.8: **Total chromatogram of fraction 6 (b) fragmentation pathway of  $m/z=255.0653$**

Fig. 5.9: (a) Chromatogram of a standard solution of 7'4' dihydroxy flavone.  
(b) Enhanced product ion and fragmentation pattern

Fig.5.10: Elemental composition of  $m/z=285.0775$  determined by MassLynx v4.1 software

Fig. 5.11: proposed fragmentation pathway of  $m/z= 285.0775$

## Chapter 6:

Fig.6.1: Inhibition of *F. solani* and *A. niger* by the crude methanol extract of *T. spicata*.

Fig. 6.2: Inhibition of *F. solani* and *A. niger* by the crude methanol extract of *T. gladiata*.

Fig.6.3: Inhibition of *A. niger* by the saponin extract and *F. solani* by the sapogenin extract of *T. spicata*.

Fig. 6.4: Inhibition of *F. solani* by (a) Toluene and ethyl acetate extract. (b) Inhibition of *F. solani* by sapogenin fraction no. 11 eluted by silica gel column chromatography.

Fig 6.5: Inhibition of *F. solani* by the compounds eluted after TLC separation

Fig 6.6: Elemental composition of  $m/z=299.1134$  determined by MassLynx v4.1 in LC-MS analysis

Fig 6.7: (a) Product ion mass spectrum obtained by ESI-QqTOF-MS/MS and (b) Mass error of the fragment ions in the product ion mass spectrum obtained by ESI-QqTOF-MS/MS

Fig. 6.8: Proposed fragmentation pathway of [(2S, 3R, 4S, 5R, 6R)-1-butyl- 3, 4, 5 trihydroxypiperidine -2yl] methyl dihydrogen phosphate from (+) ESI-IT-MS<sup>n</sup>. Accurate mass measurements of ions acquired by ESI-QqTOF-MS/MS

Fig 6.9: (a) DEPT <sup>13</sup> C NMR analysis (b) <sup>13</sup> C NMR spectrum of the purified compound

Fig 6.10: <sup>1</sup>H NMR spectra of the purified compound

Fig 6.11: Chemical structure of [(2S, 3R, 4S, 5R, 6R)-1-butyl- 3, 4, 5 trihydroxypiperidine -2yl] methyl dihydrogen phosphate

Fig 6.12 (a) Chemical structure of [(2S, 3R, 4S, 5R, 6R)-1-butyl- 3, 4, 5 trihydroxypiperidine -2yl] methyl dihydrogen phosphate and (b) Miglustat [(2R, 3R, 4R, 5S)-1-butyl-2-(hydroxymethyl)piperidine-3,4,5-triol]

## Chapter 7:

Fig. 7.1: Calibration curve of the TLC method for quantification of diosgenin

Fig. 7.2: The three dimensional image of the calibration samples

Fig. 7.3: HPTLC profiles of diosgenin samples from aerial parts of *Trigonella*

Fig. 7.4: (a) Chromatogram of *T. foenum-graecum* extract. (b) Chromatogram of *T. spicata* extract. Peak 3 is of diosgenin ( $R_f: 0.15 \pm 0.02$ ). Peaks 1-9 belong to other components present in the extract

Fig. 7.5: HPTLC profile of diosgenin samples from seeds of some *Trigonella* species