

BIBLIOGRAPHY

- Abbott, W.S. (1925): A method of comparing the effectiveness of insecticides. *J. Econ. Ent.*, **18**: 260-267.
- Adedire, C.O. and Akinneye J.O. (2004): Biological activity of tree marigold, *Tithonia diversifolia* on cowpea seed bruchid, *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Annals. Of App. Biology*, **144**(2): 185-189.
- Adiroubane, D. and Letchoumanane, S. (1998): Field efficacy of botanical extracts for controlling major insect pests of okra (*Abelmoschus esculentus*). *Ind. J. Agri. Sci*, **68**(3): 168-170.
- Agarwal, N.S. (1955): Bionomics of *Atractomorpha crenulata* Fab. *Ind. J. Ent.*, **17**(11): 230.
- Ahmed, M.; Gupta, B.K. and Bhandari, R.S. (1991): Efficacy of some plant extracts against Ailanthus Web Worm: *Atteva tebiicelle*, *Ind. J. Forestry*, **14**(1): 5-7.

- Ahmed, S.R. and Bhattacharya, A.K. (1991): Growth inhibitory effect of some plants for *Spilosoma obliqua* Walker. Ind. J. Ent., **53**(3): 453-474.
- Amicot, M.; Castella, C.; Cuany, A.; Berge, J.B. and Pauron, D. (1992): Target modification as a molecular mechanism of pyrethroid resistance in *Drosophila melanogaster*. Pesticide Biochemistry and Physiology, **44**(3): 183-190.
- Apablaza, J.U. and Norero, A.L. (1993): Captures of *Agrotis ipsilon*, *Heliothis Zae* and *Trichoplusia ni* (Lepi: noct) with synthetic pheromone trapes in the Metropolitan region (in spanish, English, Summary). Cienciae Investigation Agraria **20**: 73-81.
- Areekul, S. (1986): Toxicity tests of some insecticides against fruit flies *Dacus cucurbitae coquillett* and *Dacus tun* (Walker). Kasetsart. J. Nat. Sci., **20**(3): 274-279.
- Arya, A. (2002): Soil solarisation in integrated pest management. In 'Plant Pest Management'.

Edited by PC. Trivedi, p.136.

- Awasthi, M.D. (1985): Persistence pattern and safety evaluation of synthetic pyrethroids on egg plant. *Ind. J. Agri. Sci.*, **55**(9): 600-603.
- Axelsson, B., Lohm, U., Person, T., Nilsson, A. and Tenow, O. (1975): Energetics of larval population of *Operophtera* spp. (Lepidoptera: Geometride) in Central Sweden during a fluctuation. *Low. Zoon.* **3**: 71-84.
- Ayyangar, G.S.G. and Rao, P.J. (1989): Neem (*Azadirachital indica* A. Juss) extracts as larval repellents and ovipositional deterrents to *Spodoptera litura* (Fabr.). *Ind. J. Ent.*, **51**(2): 121-124.
- Azam, K.M. and Ali, M.H. (1965): Morphology of -I larva of *Dichocrocis punctiferalis* Guen., the Castor shoot and capsule borer with special reference to its chaetotaxy (Lepidoptera: Pyralidae). *Indian J. Ent.* **27**(4): 423-431.

- Azeezbasu, A.; Chelliah, S.; Gopalan, M. (1982): Effect of synthetic pyrethroids in the control of brinjal fruit borer, *Leucinoides orbanalis* Guen. Pesticides, **16**(1): 1-11.
- Babu, M.H., Bhattacharya, A.K. and Rathore, Y.S. (1978): Developmental behaviour of three lepidopterous pests on soyabean and green gram. Z. ang. Ent. **85**: 108-112.
- Bailey, C.G. and Singh, N.B. (1977): An energy budget for *Mamestra configurata* (Wlk.) (Lepidoptera: Noctuidae). Can. Ent. **109**: 687-693.
- Balasubramanian, G. and Balasubramanian, M. (1984): Susceptibility of different larval instars of *Spodoptera litura* F. to insecticides. Ind. J. Agri. Sci., **54**(1): 72-74.
- Bamaiyi, L.J., Ndams, I.S., Toro W.A. and Odckina S. (2007): Laboratory evaluation of Mahogany (*Khaya senegalensis* (Desv.) seed oil and seed powder for the control of *Callosobruchus maculatus* (Fab.) (Coleoptera: Bruchidae) on

- stored cowpea. J. of Entomology, **4**(3): 237-242.
- Banerjee, S.K. and Katiyar, K.N. (1984): Insecticidal control of cotton pests. Ind. J. Ent., **46**(1): 1-11.
- Basheer, S.; Vasudev, V.; Renu, R.; Guruprasad, K.P. and Harish, S.K. (1999): Toxic effects of a recently introduced carbamate pesticide, Dumet (methomyl) on *Drosophila melanogaster*. J. of Environ. Bio., **20**(2): 135-139.
- Basu, A.C. (1943): Life history and bionomics of the cauliflower pest *Prodenia litura* F. in Bengal, Sci. Cult. **10**: 420-422.
- Bathal, S.S.; Singh, D.; Dhillon, R.S.; Nayyar, K. and Singh, D. (1991): Ovicidal effects of neem oil and plant extract of *Ageratum conyzoides* Linn. on *Dysdercus koenigii* Fab. J. Insect. Sci.,

- Beck, S.D. (1956): Ibid. **49**: 552-557.
- Behera, U.K. and Satapathy, C.R. (1996): Screening of indigenous plants for their insecticidal properties against *Spodoptera litura* Fb. Insect Environ., **2**(2): 43-44.
- Benerjee, T.C. and Haque, N. (1983): Biology of *Diacrisia casignetum* kollar (Lepidoptera: Arctiidae). Indian J. Agric. Sci. **53**(5): 370-375.
- Benerjee, T.C. and Haque, N. (1984): Dry matter budgets for *Diacrisia casignetum* kollor larvae fed on sunflower leaves. J. Insect Physiol. **30**(11): 861-866.
- Bestmann, H.J.; Pietschmann, M.; Steinmeier, K. and Vostrowsky, O. (1991): Volatile constituents from *Crotalaria ochroleuca* and their effect on pest insects. Zeitschrift-fur-Naturforschung. Section C Biosciences, **46**(7-8): 579-584.
- Bhaduri, N.; Ram, S. and Patil, B.D. (1985): Evaluation of some extracts as protectants against pulse

- beetle *Calosobruchus maculatus* (Fabricius) infesting cowpea seed. J. Ent. Res., **2**: 183-187.
- Bhathal, S.S.; Singh, D.; Dillon, R.S. and Singh, D. (1994): Insecticidal activity of *Ageratum conyzoides* Linn. against *Lipaphis erysimi* (Kaltenbach). J. Insect Sci., **7**(1): 35-36.
- Bhatia, R. and Verma, A.K. (1995): Relative toxicity of insecticides against the cabbage caterpillar *Pieris brassicae* (L.) and cabbage aphid *Brevicoryne brassicae* (L.). Pest. Management Eco. Zoo., **3**(2): 93-96.
- Bhattacharya, A.K. and Pant, N.C. (1976): Studies on the host plant relationships: consumption and utilisation profile in insects. Proc. Nat. Acad. Sci. India. **46**(13): 273-301.
- Bhattacharya, A.K. Yadav, S.R. and Chaudhary, R.R.P. (1977): Biology of *Plusia orichalsia* Fabr. (Lepidoptera: Noctuidae). Sci. Cult. **43**(4): 173-174.

Bhattacharjee, N.S. and Gupta, S.L. (1971): Identity of lepidopterous pests infesting wheat crop at Delhi. Indian J. Ent. **33**(4): 435-451.

Bhumannavar B.S. and C.A. Viraktamath (2001): Larval host specificity, adult feeding and oviposition preference of the fruit piercing moth, *Othries homaena* Hubner (Lepidoptera: Noctuidae) on different Manispermaceae host plants. J. Ent. Res. **25**(3): 165-181.

Bhumannvar, B.S. (2000): Studied on fruit piercing moths (Lepidoptera: Noctuidae) species composition, biology and natural enemies. Ph.D. Thesis. University of Agricultural Science, Bangalore.

Boetal, A.M.; Fuller, W.B.; Chander, D.L.; Hovland, G.D. and Evenson, G.P. (1998): Fecundity and egg viability of northern and western corn rootworm (Coleoptera: Chrysomelidae) beetles surviving labelled and reduced soil insecticide applications. J. Eco. Entomology, **91**(1): 274-

- Borle, M.N.; Doiphode, S.M. and Deshmukh, S.D. (1979):
Efficacy of some modern synthetic insecticides
against certain important pests of sorghum.
Pesticides, **13**(8): 25-26.
- Breeland, S.G. (1958): Biological studies of the
armyworm, *Pseudaletia unipuncta* (Haworth),
in Tennessee (Lepidoptera: Noctuidae). J.
Tenn. Acad. Sci. **33**: 263- 347.
- Breuer, M. and Schmidt, G.H. (1990): Studies on the
effect of *Melia azedarach* extracts on
Spodoptera frugiperda (J.E. Smith)
(Lepidoptera: Noctuidae). Mitteilungen der
Deutschen-Gesellschaft-fur-Allgemeine und
Angewandte-Entomologie. **7**(4-6): 419-429.
- Briggs, G.G., Elliott, M., Farnham, A.W. and, Janes, N.F.
(1974): Structural aspect of the knockdown of
pyrethroids. Pestic.Sci. **5**: 643-649.
- Browning, H.W., Fedrici, B.A. and Oatman, E.R. (1982):

Occurrence of a disease caused by a Rickettsia like organism in a larval population of the cabbage looper, *Trichoplusia ni*. in southern California, environmental Entomology. **11**: 550-554.

Bull, D.L. and Pryor, N .W. (1990): *In vivo* and *in vitro* fate of fenvalerate in house flies. Pesticides Biochemistry and Physiology, **38**: 140-152.

Busbey, R.L. (1950-51): "Plant that help kill insects" in United States Department of Agriculture. The year book of Agriculture. Crop in Peace and War. Govt. Printing Office, Washington, D.C., pp.765-771.

Butter, N.S. and Surjit Singh (1996): Feeding preference of *Helicoverpa armigera* for cotton Genotypes. India J. Ent. **58**(2): 108-111.

Casid, J.E.; Grammon, D.W.; Glickman, A.H. and Lawrence, L.J. (1983): Mechanism of selective action of Pyrethroid insecticides. Ann. Rev. Pharmacol., Toxicol. **23**: 413-438.

- Charensom, K. (1992): Entomologous Insects, Department of Entomology, faculty of, Agriculture, Kasertsart University.132 P.P. (in Thai).
- Chari, M.S.; Patel, S.N.; Vaze, G.M. and Patel, L.A. (1983): Need based application of cypermethrin (Ripcord: 10 EC) for the control of bollworms on hybrid and cotton. Pesticides, **17**(3): 14-16.
- Chaudhari *et al.* (2001): Incidence of insect pests of Cabbage in relation to prevailing climatic conditions of tarai region. Indian J. Ent.**63**(4): 421-428.
- Chitra, K.C.; Janardhan, S.; Kameswar Rao, P. and Nagaiah, K. (1993): Field evaluation of certain plant products in the control of Brinjal Pest Complex. Ind. J. Ent., **55**(3): 237-240.
- Chitra, K.C.; Rao, S.J. and Rao, P.K. (1997): Efficacy of plant extracts for controlling cotton aphid (*Aphis gossypii*). Ind. J. Agric. Sci., **67**(3):

134-135.

Chlodny, J. (1967): The energetics of the development of cabbage white *Pieris brassicae* L. (Lepidoptera). Ekol. Pol. Seria A. **15**: 553-561.

Chopra, R.N. and Bhadwor, R.L. (1949): Poisonous plants of India. Scientific Monograph, ICAR, New Delhi, pp.10-12.

Chou, Y.M., Rock, G.C. and Hodgson, E. (1973): Consumption and utilization of chemically defined diets by *Argyrotaenia velutinana* and *Heliothis virescens*. Ann. Ent. Soc. Am. **66**(1): 627-632.

Consoli, R.A.G.B.; Mendes, N.M.; Pereira, J.P.; Santos, B.S. and Lamownier, M.A. (1988): Influence of some plant extracts on the survival of larvae of *Aedes fluviatilis* (Lutz) (Diptera: Culicidae) in the laboratory. Memories do Instituto Oswaldo Cruz, **83**(1): 87-93.

Cooper, M.R. and Johnson, A.W. (1984): In reference

book 161, Ministry of Agriculture, Fisheries and Food, London.

Crampton, G.C., (1921): The sclerites of the head and the mouthparts of certain immature and adult insects Ann. Ent. Soc. Am. **14**(2): 65-110.

Creamer, N.G. and K.R. Baldwin (2000): An evaluation of summer cover crops for use in vegetable production systems in north Carolina. Hort science (Print) July **35**(4): 600-603.

Creighton. C.S. Mc Fadden T.L. and Robbins. M.L. (1981): Comparative control of caterpillars on cabbage cultivary treated with *Bacillus thuringiensis*. Journal of the Georgia Entomological society **16**: 361-367.

Crumb, S.E. (1934): A classification of some noctuid larvae of the sub-family Hypernidae. Entomologica Am. **14**: 133- 196.

Dadmal, S.M.; Pawar, N.P.; Ghawde, S.M. and Kale, KB. (2001): Efficacy of plant products and

chemical insecticides against the citrus thrips, *Scritothrips citri* Moulton. Pest Management and Economic Zoology, **9**(1): 93-95.

Dandale, H.G.; Khan, K.M.; Thakar, H.S. and Borle, M.N. (1981): Comparative efficacy of synthetic pyrethroids against pod borer complex of red gram. Indian J. Ent., **43**(4): 416-419.

Dani, A., Sharma, N. and Sehagal, S.S. (1980): The method of sexing pupae and adults of Castor semilooper, *Achoea janata* (Linn). Indian J. Ent. **42**(2): 310-311.

Danks H.V. (2001): The nature of dormancy responses in insects. Acta. Soc. Zool. Bohem **65**: 169-179.

Das, B.B. (1988): Efficacy of some synthetic pyrethroids and conventional insecticidal spray against groundnut pest complex. Pesticides, **22**(1): 10-12.

Das, N.D.; Maruthi Shankar, G.R. and Biswar, K.K.

(2000): Field evaluation of Botanical and Biopesticides against pod borer, *Helicoverpa armigera* on pigeonpea. Ann. Pl. Protec. Sci., **8**(2): 233-280.

Dash, A.N.; Mukherjee, S.K. and Sontakke, B.K. (2001): Efficacy of some commercial neem formulations against major pests of rice and their safety to natural enemies. Pest Management and Economic Zoology, **9**(1): 59-64.

Dass, P., B.G. Unni., P.R. Bhattacharya and P.C. Deka (2002): Seasonal changes in food consumption and utilization pattern of semidomesticated muga silkworm, *Antheraea assama*. Westwood (Lepidoptera: Saturniidae). J. Ent. Res. **26**(4): 277-284.

David, P. and Leutan, L. (1986): Pesticides: Amounts applied and amounts reaching pests. Bioscience, **36**(2): 86-91.

De Gryse, J.J. (1915): Some modifications of the

hypopharynx in lepidopterous larvae Proc.
Ent. Soc. Wash. **17**: 173-179.

Deering M.D., Haslitt T. and Scriber J.M. (2002): Factors
affecting diapause termination in *Papilio
trailus* pupae (Lepidoptera: Papilionidae)
Holarctic Lepidoptera **8**: 43-47.

Deering M.D., Scriber J.M. (2002): Field bioassays show
hetero specific mating preference assymetry
between hybridizing North American Papilio
butterfly species (Lepidoptera: papilionidae) J.
Ethology published "on line" March 2002.

Deka, N.K.; Prasad, D. and Chand, P. (1989): Chemical
control of *Heliothis varmigera* (Hub.). Res.
Develop. Reporter, **6**(1): 130-137 .

Delabic, J.; Christian F.C. and Masson, C. (1985): Toxic
and repellent effects of cypermethrin on the
honey bee (*Apis melliferalingustica*). Pestic.
Sci., **16**(4): 409-415.

Denlinger D.L. (2002): Regulation of diapause Ann.Rev.

Entomol **43**: 93-122.

Deshmukh, P.D.; Rathore, Y.S. and Bharracharya, A.K.
(1977): Developmental behaviour of *Diacrisia obliqua* Walker (Lepidopera: Archidae) on sixteen plant species. Z. ANgew. Ent., **84**: 431-435.

Deshmukh, P.D.; Rathore, Y.S. and Bhattacharya, A.K.
(1979): Larval survival of *Diacrisia obliqua* Walker on several plant species. Indian J. Ent., **41**: 5-12.

Deshmukh, P.D.; Rathore, Y.S. and Bhattacharya, A.K.
(1976): Host range of Bihar hairy caterpillar *Diacrisia obliqua* Walker. Bull. **2**: 17-25.

Devi, Y.; Prameela and Kumar Nanda, N. (1981): Identification of cholinesterase antagonistic compound in *P. hysterophorus*. Baroda J. Nutr., **8**: 104-106.

Dhaliwal, G.S.; Pathak, M.D. and Vega, C.R. (1990): Effect of rice allochemical on insect pests,

- Predators and Plant Pathogens. J. Insect Science, **3**(2): 136-140.
- Dhandapani, N. and Balasubramanian, M. (1980): Consumption and utilization of different food plants by *Heliothis armigera* (Hubner) (Noctuidae: Lepidoptera). Entmon **5**(2): 99-103.
- Dhingra, G.B.; Chowdhry, J.B. and Sareen, P.K. (1988): Genotoxic effects of a synthetic pyrethroid insecticide on *Drosophila*. *Drosophila Inform. Serv.*, **67**: 27.
- Diaz, B; Garzo, E; Duque, M; Gonzalez P; Ferreres. A. (2002): Particle film: Effect on the mortality and development of *Trichoplusia ni* Hubner. *Boletin-de-Sanidad-vegetal, -plagas.* **28**:2, 177-183; 10 ref.
- Downey, J.C. and Allyn, A.C. Jr. (1979): Morphology and biology of the immature stages of *Leptotes cassius theonus* (Lucas) (Lepidoptera: Lycaenidae). Bull. Allyn Mus. **55**: 1-27.

Dubey, N.K.; Kishore, N.; Tripathi, N.N.; Tripathi, R.D. and Dixit, S.N. (1981): Fungitoxicity of the essential oil of *Ocimum canum* against *Aspergillus flavus* and *A. versicolor*. Indian Perfumer., **1**(2): 22-26.

Dubey, A.K. and Tripathi, N.N. (1993): Volatile antifungal activity in leaf extracts of some higher plants. National Academy Science Letters, **16**(7-8): 217-218.

Duhoon, S.S. and Banerjee, S.K. (1986): Efficacy of some synthetic pyrethroids for the control of bollworms in desi cotton (*Gossypium arboreum* Linn.). Ind. J. Entomol., **46**(2): 122-125.

Duporte, E.M. (1961): Manual of Insect morphology Reinhold, New York.

Dyar, H.G. (1890): The number of molts of lepidopterous larvae. Psyche. **5**: 420- 422.

Elliot, M. and Jams, N.F. (1978): Synthetic pyrethroids -

a new group of insecticides. Chem. Soc. Rev.,
7: 473-494.

Elsey K.D., Rabb R.L. (1970): Analysis of the seasonal mortality of the cabbage looper in North Carolina. Annals of the entomological society of America **63**: 1597-1604.

Evans, A.C. (1939a): The utilization of food by the larvae of the buff tip, *Phaleria bucephata* (Linn.) (Lepidoptera). Proc. R- Ent. Soc. Lond(A), **14**: 25-30.

Evans, A.C. (1939b): The utilization of food by certain lepidopterous larvae. Trans. R. Ent. Soc. London, **89**: 13-22.

Farooqui, T.N.A. and Singh, K.M. (1973): Salient morphological features of mature larva of bagworm moth, *Clania crameri* West. (Lepidoptera: Psychidae). Indian. J. Ent. **35**(3): 264-270.

Faweett, C.H. and Spencer, D.H. (1970): Plant

- chemotherapy with natural plant products.
Annu. Rev. Phytopathol., **8**: 403-418.
- Feinstein, L. (1952): Insecticides from plants. USDA
Washington, pp.222-229.
- Fenny, P.P. (1975): Biochemical coevaluation between
plants and their insect herbivorus.
Coevolution of Animals and Plants, pp.3-19.
- Fischer K. and Fielder K. (2001): Diamorphic growth
patterns and sex specific reaction norms in
the butterfly *Lycaena hippothoe sumadiensis*
J. Evol. Biol. **14**: 210-218.
- Fitt, G.P. and Daly, J.C. (1990): Abundance of over
wintering pupae and the spring generation of
Helicoverpa spp. (Lepidoptera: Noctuidae) in
northern new South Wales, Australia,
consequences for pest management. Journal
of Economic Entomology. **83**, 1827-1836.
- Forbes, W.T.M. (1910): A structural study of some
caterpillars. Ann. Ent. Soc. Am. **3**(1): 94-143.

- Forbes, W.T.M. (1916): On certain caterpillar homologies
J.N.Y. Ent. Soc. Am. **24**: 137-142.
- Fracker, S.B. (1915): The classification of lepidopterous
larvae. Illinois Biol. Monogr. **2**: 1-169.
- Froankel.G. (1959): The raison d'etre of secondry plant
substances. Science, **129**: 146-170.
- Fukami, H. and Nakajima, M. (1971): Rotenone and the
Rotenoids I Naturally Occurring insecticides
(eds. M. Jacobson and D.G. Gosby). pp.71-79
(New York Marcell Dekker).
- Gains, J.C. and Campbell, F.L. (1935): Ayar's rule a
related to the number of instars of the corn
ear worm *Helicthis obsoleta* Fab. collected in
the field. Ann.Ent. Soc. Am. **28**(4): 445-461.
- Ganguly, L.K.I. and Das, A.R. (1993): *In vitro* antifungal
activity of some plant extracts. Environ. Ecol.,
11: 952-954.
- Gardner, J.C.M. (1946a): On the larvae of the Noctuidae
(Lepidoptera)-I. Trans. R. Ent. Soc. London,

96: 61-72.

Gardner, J.C.M. (1946b): On the larvae of Noctuidae (Lepidoptera)-II. Trans. R. Ent. Soc. London, **96:** 237-252.

Gardner, J.C.M. (1947): On the larvae of Noctuidae (Lepidoptera)-II. Trans. R. Ent. Soc. London, **98:** 59-89.

Gautam, R.K. and Gunjan Verma (2005): Effect of thermoperiod on the regulation of diapause in *Heliothis armigera* Hubner (Lepidoptera: Noctuidae) PP. 97. VII National symposium on Indian Entomology; productivity & health. Organized by Uttar Pradesh Zoological Society.

George, G. and Patel, J.R. (1992): Mint *Mentha Spicata* - a promising botanical protectant for green gram against pulse beetle, *Callosobruchus analis* F. Indian J. of Plant Protection, **20**(1): 66-69.

Gerard, P.J. and Rub, I.D. (1991): Screening of plants and plant extracts for repellency to *Tinea dubiella*, a major New Zealand wool pest. Proc. of the 44th New Zealand Weed and Pest Control Conference (edited by Popay, A.J.). pp.205-208. Palmerston New Zealand Weed and Pest Control Society, Inc.

Gerasimov, A.M. (1935): Zur Frage der Homodynamic der Borsten von schmetterlingsranpen. Zool. Anz.**112**: 117-194.

Ghatak, S.S. and Bhusan, T.K. (1995): Evaluation on the ovicidal activity of some indigenous plant extracts on Bihar hairy caterpillar, *Spilosoma obliqua* (WK) (Arctiidae: Lepido petra). Environ. Ecol., **13**(2): 294-296.

Goel, S.C. and Kumar, A. (1981): Cephalic demarcations and chaetotaxy of larval *Diacrisia oblique* (Walk.) (Arctiidae). Uttar Pradesh J. Zool.**1**: 6-15.

Goel, S.C. and Kumar, A. (1983): Morphology and weight

loss in the pupa of *Diacrisia oblique* (Walk)
(Lepidoptera: Arctiidae). Ann. Ent. Soc.
Am. **76**: 437-440.

Gonzalez, R.H.; Buyekx, K.J. and Brader, L. (1977): In
Natural products and the protection of plants
(ed.) Marini-Bettolo, G.B. Elsevier Scientific
and Publishing Company, New York, pp.687-
707.

Grainage, M.; Ahmed, S.; Mitchell, W.C. and Hylin, J.
(1985): Plant species repeatedly possessing
pest control properties. An EWCI/VH Data
Base resource systems Institut, East-West
Center, Honolulu, Hawaii, U.S.A.

Green, M.M.; Singer, J.M.; Sutherland, D.J. and Hibben,
C.R. (1991): Larvicidal activity of *Targets*
minuta (Marigold) towards *Aedes aegypti*. J.
American Mosquito Control Association, **7**(2):
282-286.

Guddewar, M.B.; Shukla, A. and Pandey, S. (1992):
Insecticidal effect of chandani

- (*Tabernaemontana coronaria*). Pl. Protn. Bull. Faridabad, **44**(4): 32-33.
- Guddewar, M.B.; Shukla, A.; Chandra, R.; Pandey, S. and Saini, M.L. (1994): *Tabernaemontana coronaria* R. Br. (Apocynaceae), a potential source of botanical insecticides. Pl. Prot. Bull. Faridabad, **46**(1): 1-5.
- Gujar, G.T. and Mehrotra, K.N. (1983): Juvenilizing effect of azadirachtin on a noctuid moth, *Spodoptera litura* Fabr. Indian J. Exp. Biol., **21**: 292-293.
- Gunaseelan, V.N. (1998): Impact of anaerobic digestion on inhibition potential of parthenium solids. Biomass and Bioenergy **14**(2): 179-184.
- Gunjan Verma and R.K. Gautam (2005): Effect of thermoperiod on the regulation of diapause in *H. armigera* Hub. (Lepidoptera: Noctuidae). J. Ent. Res., **29**(3): 257-258.
- Gunjan Verma and R.K. Gautam (2006): Bioenergetics of *H. armigera* on *gieerarietum* plant with

different parameters on live weight basis. J. Ent. Res., **30**(40): 1-3.

Gupta, B.M.; Singh, R.; Pandey, S.N. and Singh, R. (1995): Control of gram pod borer *Heliothis armigera* (Hubner.) by reduced doses of insecticidal dusts and synthetic pyrethroids on gram. Ind. J. Entmol., **57**(2): 161-164.

Gupta, G.P. and Katiyar, K.N. (1985): Bioefficacy and economics of synthetic pyrethroids for the control of cotton bollworms. Indian J. Ent., **47**(4): 381-387.

Gurusubramanian, G. and Krishna, S.S. (1996): The effects of exposing eggs of four cotton insect pests to volatiles of *Allium sativum* (Liliaceae). Bulletin of Entomological Research, **86**(1): 29-31.

Halpern, M.E. and Morton. R.A. (1987): Reproductive and developmental defects in a Malathion-resistant, laboratory selected population of *Drosophila melanogaster*. Pesticide

Biochemistry and Physiology, **28**: 44-56.

Hansen, J.D. (2002): Effects of cold temperature treatments on the mortality of eggs and feeding larvae of the oriental fruit moth. Hart Tech. **12**: 203-205.

Haque. K.E.; Roy, Lala, P.K. and Dutta, C.P. (1984): Phytochemical investigations of Parthenium Linn. An irritating promising Weed. J.Indian Chern. Soc., **61**: 92.

Heinrich, C. (1916): On the taxonomic value of some larval characters in the Lepidoptera Proc. Ent. Soc. Wash. **18**: 154-164.

Hifnawy, M.S.; Abdel-Wahab, S.M.; El-Hawary, S.S. and Karawaya, M.S. (1990): Study of essential oil of *Artemisia monosperma* and its larvicidal effect. International J. Crude Drug Research, **28**(4): 247-251.

Hinton, H.E. (1946): On the homology and nomenclature of the setae of lepidopterous larvae with some

- notes on the phylogeny of the Lepidoptera.
Trans. R. Ent. Soc. London. **97**: 1-37.
- Hinton, H.E. (1963): The origin and function of pupal stage. Proc. B. Ent. Soc. Lond. (A), **38**: 77-85.
- Hiremath, I.G. and Ahn, Y.J. (1997): Parthenium as a source of pesticide. In First International Conference on Parthenium Management, Dharwad, India, Oct. 6-8, pp.18-89.
- Hossain, M.l.; Ameen, M. and Ahmed, A.K.M.R. (1995): Effect of two pyrethroid insecticides against *Culex quinquefasciatus* Say larvae in Dhaka city. Bangladesh Journal of Zoology .. **23**(2): 187-192.
- House H.L. (1969): Effect of different proportions of nutrient on insects. Ent. Exp. And Appl. 12: 651-669.
- Howse P.E. (1994): Laboratory evaluation of toxic and repellent properties of the Pithraj tree, *Appanamixis polystachya* Wall and Parker,

- against *Sitophilus oryzae* (L). Int. J. of Pest Management. **40**(3): 274-279.
- Hughes, R.D. (1960): Induction of diapause in *Erioisehia brassica* Biouche (Dept. Anthomyiidae). J. Exp. Biol., **37**: 218-223.
- Imms, A.D. (1973): A general Text Book of Entomology (revised) x+ 886 E.L.B.S. London.
- Ishman, Murry B. and Rodriques, Eloy, (1983): Larval growth inhibitory from species of Parthenium (Asteraceae). Phytochemistry, **22**: 2209-2213.
- Islam, N.; Bhuiyah, M.I.M.; Begum, A. and Karim, M.A. (1990): Field evaluation of foliar insecticides for the control of the mustard aphid, *Lipaphis erysimi* (Kalt). Bangladesh Journal of Zoology, **18**: 185-188.
- Issac, P.V. and Rao, K.V.A. (1941): A key for the identification of the known lepidopterous borers of sugarcane in India based on morphological characters. Indian J. Agric.

Sci. **11**: 795-803.

Jackai, L.E.N.; Inang, E.E. and Nwobi, P. (1992): The potential for controlling post flowering pests of cowpea, *Vigna unguiculata* Walp. using neem, *Azadirachta indica*. A Jurs. Tropical Pest Management, **38**(1): 56-60.

Jackai, L.E.N. and Oyediran, I.O. (1991): The potential of neem *Azadirachta indica* A. Juss. For controlling post flowering pests of cowpea, *Vigna unguiculata* Walp. The pod borer, *Maruca testulalis*. Proceedings of the second International Conference on Tropical Entomology, July 31-August 4, 1989, held in Nairobi, Kenya.

Jackbson, M. (1975): Insecticides from plants. A Review of Literature (1954-1971), USDA Agric. Hand book, Govt. Printing Office Washington, D.C. 461-1975.

Jacob, S. and Sheila, M.K. (1996): Bioefficiency of few synthetic pyrethroids against *Aphis gossypii*

on Okra. Indian J. Ent., **58**(2): 112-114.

Jaglan, M.S.; Khokar, K.S.; Malik, M.S. and Singh, Ram (1997): Evaluation of neem (*Azadirachta indica* A. Juss) extracts against American bollworm, *Helicoverpa armigera* (Hubner). J. Agric. Food Chemistry, **45**(8): 3362-3268.

Jaiswal, A.P. and Saini, R.K. (1981): Effect of some synthetic pyrethroids on pink bollworm incidence and yield of cotton. Pesticides, **19**(1): 33-35.

Jaques, R.P. (1988): Field testson control the imported cabbage worm (Lepidoptera: Pieridane) and the cabbage looper, the Canadian Entomologist 120, 575-580.

Jaques, R.P. and Laing J.E. (1984): *Artogeia rapae* L), imported cabbage worm (Lepidoptera: pieridae) *Trichoplusia ni* (Hubner), Cabbage looper (Lepidoptera: Noctuidae) and plutella xylostella a (L), diamond back moth (Lepidoptera: Plutellidae). In Biological control

programmes Against Insects and Weeds in Canada. 1969-1980.

Jaya Varma and Dubey, N.K. (1999): Prospectives of botanical and microbial products as pesticides of tomorrow. *Current Science*, **76**(2): 172-179.

Jost. D.J; Pitre H.N. (2002): Soyabean looper and cabbage looper (Lepidoptera: Noctuidae) populations in cotton and soyabean cropping systems in Mississippi. *Journal of Entomological-science*.**37**:3, 227-235.

Justin, C.G.L.; Rabindra,R.J. and Jayaraj, S. (1995): Combined efficacy of *Bacillus thuringiensis* Berliner and some plant extracts against *Plutella xylostella* (L.) larvae. *Pest Management and Economic Zoology*, **3**(2): 75-77.

Kallapur, V.L. Majumder, C.; Roe, R.M. (1996): *In vivo* and *in vitro*-tissue specific metabolism of Juvenile hormone during the last stadium of

the cabbage looper, *Trichoplusia ni*. Journal of Insect Physiology **42**(2): 181-190.

Kalushikov P., Hodkova M., Nedved O. and Hodek I. (2001): Effect of thermoperiodon diapause intensity in *Pyrrhocoridae apterus* (Heteroptera: Pyrrhocoridae). J. Insect Physical **47**: 55-61.

Kamal, R.; Mangla, M.; Prakash, D.; Sharma, G.L. and Srivastava, R.C. (1988): Relative toxicity of natural pyrethrins against *Mesocyclops leukartis* L. the carrier of dracunculiasis. Insect Sci. and its applications. **9**(4): 437-440.

Karthikeyan, K., Jacob Sosamma, Purushaman, S.M., Revi Smitha (2007): Efficacy of plant products and synthetic insecticides against rice blue beetle *Leptispa pygmaea* Baly (Chrysomelidae: Coleoptera). Ind. J. Crop. Sci., **2**: 73-80.

Karunasri, K., M.V. Sailaja, Et. Al. (1998): Effect of speical association of pearl millet (*Pennisetum glaucum*) and ground hut (*Arachis Hypogaea*)

on resource utilization and economic yield,
Indian Journal of Agronomy **43**(1): 33-37.

Kasting, R. and Mc Ginnis, A.J. (1959): Nutrition of the
pale western cutworm, *Agrotis orthogonia*
Morr.(Lepidoptera: Noctuidae)-11 Dry matter
and nitrogen economy of larvae fed on sprout
of a hard red spring and a durum wheat. Can.
Zool.**37**: 713-720.

Kasting, R. and Me Ginnis, A.J. (1959): Nutrition of the
pale western cutworm, *Agrotis* Ent. Soc.
Am.**57**: 98-102.

Kasting, R. and Me Ginnis, A.J. (1962): Quantitative
relationship between consumption and
excretion of dry matter by larva of pale
western cutworm, *Agrotis orthogonia*
Morr.(Lepidoptera: Noctuidae). Can Ent.
Zool.**94**: 441-443.

Kayitara, J. and Ntezurubanea, L. (1991): Evaluation of
the toxicity and repellent effect of certain
plants from Rwanda against the bean

bruchids. *Acanthoscelides obtectus* say and *Zabrotes subfasciantus* Boheman. Insect Science and its application. 1991, **12**: 5-6,695-697. In proceeding of the AAIS and the Entomological Society of Zambia Annual General and Scientific Meetings, Lusaka, Zambia, 4-8 Dec., (1989):

Kelly, T.J. and Cowdron (1990): Total and specific ecdysteriods in the haemolymph of *Trichoplusia ni* (Lepidoptera: Noctuidae) and its, parasite, *Euplectrus plathypenae* (Hymenoptera: Eulophidae). Journal of Insect Physiology **36**: 463-470.

Kenneth A, Sarensen (1966): Vegetable Insect Pest Management North Carolina cooperative Extention Service North Carolina State University **5**: (1-5).

Kenny, G.J. and Chapman, R.B. (1988): Effect of an intercrop on the insect pests, yeild and quality of cabbage. New Zealand and Journal

of Experimental Agriculture 16, 67-72.

Khanam, L.A.M.; Talukdar, D. and Khan, A.R. (1990):
Insecticidal property of some indigenous
plants against *Tribolium confusum* Duval
(Coleoptera: Tenebrionidae). Bangladesh J. of
Zoology, **18**(2): 253-258.

Kogan, M. and Cope, D. (1974): Feeding and nutrition of
insects associated with soybeans 3. Food
intake, utilization and growth in the soybean
looper, *Pseudoplusia includes* Ann. Entomol
Soc. Am. **67**(1): 66-72.

Koul, O. (1999): Insect growth regulating and antifeedant
effects of neem extracts and azadirachtin on
two aphid species of ornamental plants. J.
Biosciences, **24**(1): 85-90.

Kulkarni, N.; Joshi, K.C. and Gupta, B.N. (1997):
Antifeedant property of *Lantana camara* Var.
aculeata and *Aloe vera* leaves against teak
Skeletonizer, *Eutectona machaeralis* Walk.
Lepidoptera: Pyralida. Entomol., **22**(1): 61-65.

- Kulkarni, S.N. (1981): Comparative efficacy of two synthetic pyrethroids in control of cottonbolloworms. *Cotton Dev.*, **10**(3-4): 15-16.
- Kumar, A. (1983): On the structure of larval head appandages of *Diacrisia oblique* (Wlk.) (Lepidoptera Arctiidae). *Uttar Pradesh J. Zool.* **3**: 35-41.
- Kumar, S. and Verma, L.R. (1980): A note on the sexual diamorphisrn in the pupae of cutworm *Agrotis segetum* Schiffer Mueller. *Indian J. Ent.* **42**(2): 301-303.
- Kumar, V. and Goel, S.C. (1988): Salient features of cephalic appandages of the last instar caterpillar of *Plusia orichalcea* Fabr. (Noctuidae). *Environ. and Ecol.* **61**(1): 79-84.
- Kumari, K. and Singh, S.N. (1998): Evaluation of efficacy of botanical as insecticides against pulse beetle (*Callosobruchus chinensis* L.). *J. Phytopathol.*, **50**(2): 235-241.

Lale, N.E.S. (1992): A laboratory study of the comparative toxicity of products from three species to the maize Weevil. *Post Harvest Biology and Technol.*, **2**(1): 61-64.

Landolt peter, J. Molina oscar, H. (1955): Host finding by cabbage looper Moths (Lepidoptera: Noctuidae): Learning host odor upon contact with host foliage. IABBB Research Laboratory 1700 S. W. 23rd Drive GAINES VILLE (1) 10-31.

Landolt, P.J. (1995): Altration of female cabbage looper moths (Lepidoptera: Noctuidae), to males in the field, *Florida Entomologist* **78**: 96-100.

Landolt, P.J., Molina, O.H., Health, R.R. Ward K., Dueban, B.D. and Miller, J.G. (1996): Starvation of cabbage looper moths (Lepidoptera: Noctuidae) increases attraction to male pheromone. *Annals of the Entomological Society of America* **89**: 459-465.

- Latheef M.A. and Ortiz J.H. (1983b): Influence of companion plants on oviposition of imparted cabbage Warm Pieris rapae (Lepidoptera: Pieridae), and cabbage lopper, *Trichoplusia ni* (Lepidoptera: Noctuidad), on callard plants. The Canadian Entomologist **115**: 1529-1531.
- Latheef, M.A. and Harcourt, O.J. (1972): Quantitative study of food consumption assimilation and growth in *Leptinotassa dacemlineata* (Coleoptera: chrysomalidae) on two host plants Can. Ent. **104**: 1271-1276.
- Latheef, M.A. and Trwin, R.D. (1983): Seasonal abundance and parasitism of lepidopterous larval on *Brassica green* in Virginia. Journal of the Georgia Entomological Society **18**: 164-168.
- Lee, H.L. and Winta, R (1993): Laboratory and field evaluation of permethrin in against *Aeded* (Stegmoyia) *albopictus* Skuse larvae. Mosquito Borne Diseases Bulletin, **10**(3): 77-82.

Lefroy, H.M. (1909): Indian insect life xii + 786. Today & Tomorrow's, New Delhi.

Lundgren, L. (1975): Natural plant chemicals acting as oviposition deterrents on cabbage butterflies *Pieris brassicae* (L), *P. rapae* (L.) and *P. nazi* (L.). Zoologica Scripta 4, 253-258.

Macedo, M.E.; Consoli, R.A.G.R.; Gandhi, T.S.M.; Anjos, A.M.G. dos, Oliveira, A.B. de.; Mendes, N.M.; Quiroz, R.O.; Zani, C.L.; Dos Anjos, A.M.G. and De Oliveira, A.B. (1997): Screening of Asteraceae (compositae) Plant extract for larvicidal activity against *Aedes fluviatilis*. Memórias-do-Instituto Oswaldo Cruz., **92**(4): 565-570.

Mackey, A.P. (1978): Growth and bioenergetic of the moth, *Cyclophragma leucosticte*. Grunberg. Oecologia (Berl), **32**(3): 367- 376.

Mahadevan, A. (1982): Biochemical aspects of disease resistance, Part-I. Performed inhibitory substances prohibitions. Today and

Tomorrow's Printers and publishers, New Delhi, India, p.425.

Maheshwari, H.K. and Dwivedi, S.C. (1995): Laboratory evolution of some plant extracts against the pulse beetle *Callosobruchus chinensis* (L.) on cowpea. Pest Management and Economic Zoology, **3**(2): 107-109.

Maheshwari, H.K. and Dwivedi, S.C. (1996): Evaluation of botanicals for the management of *Tribolium castaneum* (Coleoptera: Tenebrionidae). Insect Environ., **2**(3): 72-73.

Maldonad, O. Errma, Mendoza, Graciela, Cardenas, Jorge and Ortega, Alferdo (1985): Susquiterpene lactones from *Parthenium tementosum*. phytochemistry, **24**: 2981-2985.

Malik, V.S. and Chaudhary, J.P. (1980): Biology of gram cutworm, *Agrotis ipsilon* Hubn. in the Haryana state. Bull. Ent.**21**: 70-78.

Mani, M.S. (1980): General Entomology. Oxford and IBH

publishing Co. New Delhi.

Marini Bittolo, G.B. (ed.) (1977): In natural products and the protection of plants, Elsevier Scientific Publishing Company, New York, pp.5-16.

Mathavan, S. and Bhaskaran, R. (1975): Food selection and utilization in danid butter fly. *Oecologia (Berl.)* **18**: 55-65.

Mathavan, S. and Pandian T.J. (1974): Use of faecal weight as an indicator of food consumption in some lepidopteron. *Oecologia (Berl.)*, **15**: 177-185.

Mathur, R.N. (1959): Immature stages of Indian Lepidoptera No. 12, Pyralidae. Sub-family Pyraustinae. *Indian for. Rec.* **9**: 183-210.

Mathur, Y.K. and Upadhyaya, K.D. (1980): Evaluation and economics of some modern insecticides against mustard aphid *Lipaphis erysimi* Kalt. *Pesticides*, **14**: 14-15.

Matsui, M. and Yamamoto, I. (1971): "Pyrethroids" In

naturally occurring insecticides (eds. Jacobson, M. and Croshyl, D.G., 3-70), New York: Marcell Dekker.

Maxwell, G.F. and Jennings, R.P. (1980): Breeding plants resistance to insects. New York, Ghicheter, Brisbane, Toronoto, John Wiley & Sons, p.683.

Mc Donald, S. (1981): Laboratory evaluation of new insecticides for control of red backed cutworm larvae. J. Econ. Entomol. **74**(5): 593-596.

Mc Pherson, R.M. (1993): Parasitism of *Pseudoplusia includence* (Lepidoptera: Noctuidae) by *Copidosoma truncatellum* (Hymenoptera: Encytriidae) using various laboratory procedures Journal of Entomological science **28**: 331-334.

Mehta, P.K.; Vaidya, D.N. and Kashyap, N.P. (1996): Effect of plant extracts on *Pieiiis brassicae* L. Insect Environ., **2**(3): 95-96.

- Migula, P. (1975): Energy flow through populations of satin moth *Leucoma salicis* (L.) Acta Biol. Cracov. Ser. Zool. **18**: 229-255.
- Mishra, D.N.; Mishra, A.K.; Tiwari, R. and Upadhyaya, P.S. (1989): Toxicity of some essential oil against pulse beetle, *Callosobruchus maculatus* (F.). Indian Perfumes, **33**(2): 137-141.
- Mishra, H.P. (1993): Evaluation of some synthetic insecticides and oils against mustard aphid *Lipaphis erysimi* Kalténbach. Indian Journal of Entomology, **55**: 58-61.
- Mohan, N.J. and Prasad, V.G. (1984): Role of synthetic pyrethroids in the control of brinjal pests. Ind. J. Entmol., **46**(2): 179-182.
- Mohite, A.S., D.B. Tenbhare and S.P. markar (2004): Biology and behaviour of developing stages of fruit sucking moth othries meterna Linn. (Lepidoptera: Noctuidae). J. ent. Res. **28**(1): 37-45.

- Mosher, E. (1916): A classification of the Lepidoptera based on charactes of the pupa Bull. Illinois St. Lab Nat. Hist. **12**: 14-159.
- Mukerji, M.K. and Guppi, J.C. (1970): Quantitative study of food consumption and growth in *Pseudoletia unipuncta* (Lepidoptera Noctuidae) Can Ent. **102**: 1179-1188.
- Mukerji, S. and Singh, H. (1951): Studies on the chaetotaxy of larvae of *Plusia species* (Lepidoptera: Phalaenidae). Proc. R. Ent. Soc. Lond. **20**(8): 15-24.
- Mundiwale, S.K.; Patel, J.D.; Men, U.B. and Khandalkar, H.G. (1989): Efficacy of synthetic pyrethroids against Sorghum pests. Indian J. Ent., **51**(2): 172-176 .
- Nanda Kumar.; Devi, P.V.; Prameela, Y.; Swami, K.S. and Majumdar, S.K. (1980): Insecticidal properties of Parthenium hysterophorus extracts. Comp. Physiol. Ecol., **5**: 264-267.

- Nasu. Y; Tanaka, H; Shibao. M. (2003): New lepidopterous pests attacking eggplant and cabbage in Osaka. Proceedings of the Kansai plant protection Society. No.45, 65-66; 7 ref.
- Neunzig, H.H. (1964): The eggs and early instar larvae of *Heliothis Zea* and *Heliothis vire-scens* (Lepidoptera: Noctuidae) Ann. Ent., **17**: 85-99.
- Nimbalkar, R.B. and Ajri, D.S. (1981): Efficacy of synthetic pyrethroids and two newer compounds against brinjal shoot and fruit borer, *Leucinodes orbonalis* Guen. Ind.J. Entmol. **43**(2): 202-204.
- Novranjan Pal, A.V., Dass, R. and Prasad, B. (1979): Sex determination of pupae of *Heliothis armigera* Hub. On gram. Ind. J. Ent. 41(3): 285-286.
- Oard, P.K.; Patel, J.R.; Ratanpura. H.C. and Shah, B.R. (1991): Evaluation of some insecticides for the relative efficacy against *Lipaphis erysimi* Kalt. and *Helicoverpa armigera* Hubner on cabbage. Ind. J. Plant Protect., **19**: 191-193.

- Oji, O. (1991): Use of piper guineense in the protection of stored *Zea mays* against the maize weevil. *Fitoterapia*, **62**(2): 179-182.
- Olaifa, J.I.; Matsumura, F.; Zeevaarat, J.A.D.; Mullin, C.A. and Charalambous, P. (1991): Lethal amounts of ricinine in green peach aphids (*Myzus persicae*) (Sulzer) fed on castor bean plants. *Plant Science Limerick*, **73**(2): 253-256.
- Orner, A.D. and Leigh, T.F. (1995): Sublethal effects of acephate and biphenate on fecundity, longevity and egg viability in the greenhouse whitefly (Hom. Aleyrodidae). *Journal of Applied Entomology*, **119**(2): 119-122.
- Oudhia, P. (1999): Effect of botanicals on the hatchability of Blumea leaf beetle egg. *Insect Environ.*, **4**(4): 154.
- Pal, R.K.; Tripathi, R.A. and Prasad. (1996): Relative toxicity of certain plant extracts to Khapra beetle, *Trogoderma granarium*. *Annals of Plant*

Protection Science, **4**(1): 35-37.

Pande, V.D.; Ghosh, D. and Guha, S. (1987): Toxicity of leaf extract of *Ageratum conyzoides* L. (Compositae) to red pumpkin beetle *Raphidopalpa foveicollis* (Lucas) (Coleoptera: Chrysomelidae) in Tripura. Ind. Agriculturist, **31**(14): 251-255.

Pandian, R.S.; Kalpana, M. and Manoharan, A.C. (1995): Susceptibility of the larvae of *Culex quinquefasciatus* say to different insecticides. Geobios (Jodhpur), **22**(4): 200-203.

Pao, M.S. and Patel, R.C. (1973): Biology and control of Okra semilooper, *Anomis flava* Fabr. (Noctuidae: Lepidoptera) on Okra. Indian J. Ent. **35**(3): 198-205.

Parihar, S.B.S.; Thapliyel, P. and Ramkishore (1999): Effect of botanicals of peach aphid *Myzus persical* (Sulzer). Insect Environ., **5**(1): 9-10.

Parmar, B.S. and Devkumari, C. (eds.) (1993): In

botanical and biopesticides, West Vile Publishing House, New Delhi, pp.197-199.

Patel, C.B. and Patel, R.C. (1973): Biology of *spodoptera exigna* (Hub) and the influence of larval diet. *Andhra agric, J.* **20**(1-2): 18-29.

Patel, J.R.; Jyani,D.B.; Patel, C.C. and Broad, P.K. (1993): Insecticidal property of some plants and their available formulations against *Amsacta moorei* Bulter. Botanical pesticides in integrated pest management. Rajmundry, India: Indian Society of Tobacco Science, pp.332-339.

Patel. J.J.; Patel, N.C.; Jayanti, D.B.; Patel, J.R and Patel B.D. (1997): Bioefficacy of synthetic and botanical insecticides for controlling pod borer (*Helicoverpa armigera*) and podfly (*Mananogromyza obtusa*) infesting vegetable purpose pigeonpea (*Cajanus cajan*). *Ind. J. of Agric. Sci.*, **67**(3): 117-119.

Patil, R.K.; Rayar, S.G.; Basappa, H.; Hiremath, I.J. and

- Patil, B.R. (1997): Insecticidal property of indigenous plants against *Dactynotus carthamii*, H.R.L. and its predator, *Chrysoperla carnea* L. J. Oilseeds Res., **14**(1): 71-73.
- Patil, T.M. and Hedge, B.A. (1988): Isolation and purification of resquiterpene lactone from the leaves of *Parthenium hysterophorus* L. Its allelopathic and cytotoxic effects, Curr. Sci., **57**: 1178-1181.
- Peterson, A. (1962): Larvae of insects (Part I). Columbus, Ohio.
- Pilcher, C.D., M.E. Rice, J.J. Obrycki and L.C. Lewis. (1997): Field and laboratory evaluations of *Bacillus thuringiensis* corn on secondary Lepidopteron Pests (Lepidoptera: Noctuidae) J. Econ. Entomol. **90**(2): 669-678.
- Piotrowska, S.B. and Piotrowski, W. (1997): The reaction of *Trichoderma* spp. to synthetic and natural pesticide compounds. B. Plant Extracts.

Roezniki Nauk-Rolniezych-Seria-E-ochrona-Roslin, **26**(1-2): 103-108.

Potenza, M.R.; Rossi, C.E. and Calofion, M.H. (1987): Use of the sunflower (*Helianthus annuus* L.) plant extract on the control of the bean hopper (*Empoasea krameri*) and *Diabrotica speciosa*. Germ. Ecosystema., **12**: 114-118.

Potter, C. (1935): An account of the constitution and use of and atomized white oil Pyrethrin fluid to control ploidia interpunctella Hb. and Ephestia elutella Hb. in ware houses, Ann. Appl. Biol. **4**: 769-805.

Pradhan, S., Singh S.R., Bhatia, S.K., Kundu G.G., Pradhan K.G. and Gupta M. (1972): Resistance to two major stored grain pests in world collection of what. Res. Bulletin New Series No. 1. I.A.R.I., New Delhi, 69pp.

Prakash, A. and Rao, Jagdismaii (1987): Use of chemicals as grain protectants in Rice storage ecosystem and its consequences. Bulletin of

Grain Technology, **25**(1): 65-69.

Prasad, J. and Bhattacharya, A.K. (1975): Growth and development of *spodoptera littoralis* (Bois) (Lepidoptera: Noctuidae) on several plants. Ang. Ent. **79**: 34-48.

Prasad, S.K. (1959): The association between the different pests on cabbage. Ibid, **21**(3): 206-209.

Printz, Y.A.I. (1925): The biology of *Heliothis obsoleta* F. and its control Notes App. Sci. Dept. Tiffis Bot. Gdns. Part 2, Separate 54 PP.

Puttarudriah, M. and Maheshwariah, B.M. (1956): Some observation on the life history and habits of *Anomis flava* (Fab.) (Noctuidae: Lepidoptera). Indian Colt. Grow Rew. **12**: 337-341.

Quarrie (1992): J. Earth Summit, London: The Regency Press Corporation.

Rahan, M.M. and Yadav, T.D. (1987): Efficacy of deltamethrin, cypermethrin, permethrin and

fenvalerate dusts against developmental stages of *Callosobruchus maculatus* Fab. and *C. chinensis* Linn. Indian J. Ent., **49**(3): 387-391.

Rajapakse, R.H.S. and Jayasena, K.W. (1991): Plant resistance and a biopesticides from lemongrass for suppressing *Spodoptera litura* in peanut. AgriculturalInternational, **43**(6): 166-167.

Rajasekaran, T.; Ravishankar, G.A.; Rajendra, L. and Venkataraman, L.V. (1991): Bioefficacy of pyretherins extracted from callus tissues of *Chrysanthemum cinerariaefolium*. Pyrethrum Pest., **18**(2): 52-54.

Rajendra, B. and Gopalan, N. (1979): Note on the insecticidal properties of certain plant extracts. Indian J. Agric. Sci., **49**: 295-297.

Rajeswari, E. and Mariappan, V. (1992): Effect of plant extracts on in vitro growth of rice blast (B₁) Pathogen *Pyricularia oryzae*. Int. Rice Res.

News1, **17**: 24.

Rajkumar, E.D.M. and Nandkumar, N.V. (1984): Effect of *P. hysterothorus* leaf extract and par-axon on cock roach head and gizzaha cholinesterase inhibition. *Geobios*, **11**: 11-16.

Ramasubramanian G.V. and Sundara Babu P.C. (1998): Effect of host plants on some biological aspects of spotted podborer, *Maruca testulalis*. (Lepidoptera: Pyralidae). *Indian J. Agric. Sci.*58, 618-620.

Ramesh K. and G.P. Gupta. (2004): Impact of abiotic factors on the incidence of major insect pest of Okra *Abelmoschus esculentus* under Delhi conditions *Shashpa*. **11**(2): 129-133.

Rani, R.D. and Osmani, Z. (1982): Toxicity and repellency of certain synthetic pyrethroids against *Musca domestica* (L.) and *Dysdercus cingulatus* (F.). *Pesticides*, **16**(9): 7-9.

Rani, S.; Tabassum, R.; Naqvi, S.N.H.; Ahmad, I.; Sultan,

- B. and Khan, M.F. (1997): Comparative toxicity of different formulations of cypermethrin, deltamethrin and permethrin against *Drosophila melanogaster* (K.U. Strain). J. Environ. Biol., **18**(4): 343-349.
- Rao, P.K.; Aleem, M.A. and Chitra, K.C. (1993): Efficacy of some botanicals and ash against pulse beetle, *Callosobruchus chinensis* (L.). Botanical pesticides in integrated pest management. In National Symposium, 21-22 Jan. 1990, Rajmundry, India, p.282-287.
- Rao, V.R.; Chitra, K.C. and Kameswara Rao, P. (1989): Relative toxicity of synthetic pyrethroids to *Henosepilachna vigintioctopunctata* (Fabricus). Indian J. Ent., **51**(1): 51-54.
- Rath, L.K.; Sontakke, B.K.; Jena, S.N. and Phurua, S. (1995): Bioefficacy of some insecticides against the mustard aphid *Lipaphis erysimi* Kalt. Pest Management and Economic Zoology, 3 (2): 115-117.

- Ravindranath, K. and Pillai, K.S. (1986): Control of fruit fly of bitter melon using synthetic pyrethroids. *Entomol.*, **11**(4): 269-272.
- Reddy, G.R.; Singh, M.R.; Madhusudhana, L. and Sreenivasulu (1994): Differential effects of synthetic pyrethroids on Na⁺, K⁺, ATPase in the cockroach, *Periplaneta americana*. *Ad. Bios.*, **13**(11): 49-56.
- Reddy, G.V.P. and Urs, K.C.D. (1989): Effects of *Tribulus terrestris* Linn. on the development of potato tuber moth, *Phytophthora operculella* Z. (Lepidoptera Gelechiidae). *Current Science*, **458**(4): 212-213.
- Reddy, M.V. (1983): The consumption and utilization of the castor leaves by the larvae of *Philosamia ricini* Hutt. (Lepidoptera: Saturniidae). *Proc. Syn. Ins. Ecol. and Resource Manage.* PP. 215-225.
- Reddy, M.V. and Alfred, J.R.B. (1979): Utilization of castor leaves by last instar larvae of

- silkworm, *Philosamia ricini* Hutt. (Lepidoptera: Saturniidae). *India Biologist*. **11**: 36-40.
- Reddy, P.V.R.; Chitra, K.C. and Rao, P.K. (1990): Field evaluation of certain plant extracts for the control of brinjal spotted leaf beetle, *Honosepilachna vigintioctopunctata*, *J. Insect. Sci.*, **3**(2): 194-195.
- Riaz, M.; Vasudev, V. and Krishnamurthy, N.B. (1990): Response of *Drosophila* to carbamate pesticide Furadan. *Trends Eco. Toxicol.*, pp.179-184.
- Ripley, L.B. (1923): The external morphology and embryology of noctuid larvae. 111, *Illinois Biols. Monor.* **8**: 1-169.
- Sagar, P. (1985): Comparative efficacy of synthetic pyrethroids and other insecticides against *Heliothis armigera* (Hubn.) and *Pulsia orichalcia* Fab. Pests of Japanese mint in the Punjab. *Indian J. Ent.*, **47**(2): 183-185.

- Sahayaraj, K. and Sekar, R. (1996): Efficacy of plant extract against tobacco caterpillar larvae in groundnut. *International Arachis, Newsletter*, No. 16, p.38.
- Salam, J.S.; Khuman, M.S. and Singh, M.P. (1995): The oil of *Parkia roxburghii*. G. Don, a potential insecticide. *Current Science*, **68**(5): 502.
- Santhi, M.; Krishnamurthy, M.M.; Mayuravalli, V.V.L. and Reddy, G.P.V. (1993): Effect of certain synthetic pyrethroids on the biology of pulse beetle, *Collosbruchus maculatus* Fab. and lesser grain borer, *Rhizopertha dominica* Fab. *Indian J. Ent.*, **5**(1): 94-98.
- Sarkar, G.M.; Banerjee, R.D.; Chatterjee, M.L.; Dutta, S. and Ghosh, A. (1995): Antimicrobial and insecticidal activity of an aquatic alga *Chara zeylanica* Klein exwild. *Int. Environ. Studies*, **48**(1): 29-39.
- Schroeder, L.A. (1972): Energy budget of the cecropia (Lepidoptera: Saturnidae) fed lilac leaves.

Ann. Ent. Soc. Am. **65**: 367-372.

Schroeder, L.A. (1973): Energy budget of the larvae of the moth *Pachysphinx modesta*. *Oikos*, **24**: 278-281.

Schroeder, L.A. (1971): Energy budget of larvae of *Hyalophora cercopia* (Lepidoptera) fed *Acer negundo*. *Oikos*, **22**: 256-259.

Scriber, J.M. and Slansky, F.N. (1981): The nutritional ecology of immature insects. *Ann. Rev. Ent.*, **26**: 183-211.

Secoy, D.M. and Smith, A.E. (1983): Use of plants in control of agricultural and domestic pests. *Econ. Bot.*, **37**(1): 28-57.

Sehgal, V.K. and Ujagir, R. (1990): Effect of synthetic pyrethroids, neem extracts (from *Azadirachta indica* seed Kernels) and other insecticides for the control of pod damage by *Helicoverpa armigera* (Hubner) on chickpea and pod damage-yield relationship at Pantnagar in

- Northern India. Crop. Protection, **9**(1): 29-32.
- Sellammal, N.; Parameswarn,S. and Balasubramanian, M. (1979): Efficacy of five different synthetic pyrethroids in control of bollwonns and aphid on cotton. Pesticides, **13**(10): 15-17.
- Sharma H.C. and B.A. Franzmann (2000): Biology of the legume pod Borer, *Maruca vitrate* (Fabricus) and its damage to Pigeonpea and Adzuki Bean. Insect Sci. Applic Vol. **20** No.2, P.P. 99-108.
- Sharma, B. and Tara, J.S. (1988): Comparison of consumption and utilization of Mulberry leaves in two moths, *Spodoptera liture* (F.) and *Diacrisia oblique* walk. Indian J. Ent. **50**(3): 336-342.
- Sharma, H.K. and Prasad, D. (1995): Potentialities of a few botanicals in management of root-knot, *Meloidogyne incognita* on Soybean. Annals Agric. Research, **16**(4): 473-476.

Sharma, R.N. and Joshi, V.N. (1977): Allomonic principles in *Parthenium hysterophorus* potential as insect control agent and role in the seed resistance to serious degradation Part II. The biological activity of *Parthenium* on insects. *Biovigyanam*, **3**: 225-231.

Sharma, R.N.; Gupta, A.S.; Patwardhan, S.A.; Hebbalkar, D.S.; Tare, U. and Bhonde, S.B. (1992): Bioactivity of Lamiaceae (Labiatae) plants against insect. *Ind. J. Exptl. Biol.*, **30**(3): 244-246.

Sharma, S.P.; Saini, M.L. and Goel, S.C. (1989): Effectiveness of synthetic pyrethroids against *Heliothis armigera* Hubner. on gram. *Ind. J. Plt. Prot.*, **17**(1): 59-62.

Singh, B. (1951): Immature stages of Indian Lepidoptera No. 8 Geometridae. *Indian for Rec.* **8**(7): 67-158.

Singh, B.B., K. Y. Mai, *et al.* (1999): Relative drought tolerance of major rainfed crops of the semi-

arid tropics. Indian journal of genetics and plant breeding. Nov. **59**(4): 437-444.

Singh, D.S. and Singh, J.P. (1997): Evaluation of susceptibility level of synthetic pyrethroids and non-pyrethroids against larvae of *Spodoptera litura*. Indian J. Ent., **59**(3): 245-247.

Singh, D.S. and Singh, J.P. (2000): Status of pyrethroid and nonpyrethroid insecticides to the larva of Bihar caterpillar, *Spilarctia obliqua*. Indian J. Entomol., **62**(2): 141-145.

Singh, D.S. and Sircar, P. (1986): Evaluation of synthetic pyrethroids for inhibitory effect on the hatching of eggs of *Earias vittella* Fab. Indian J. Ent., **48**(1): 101-103.

Singh, D.S. and Sircar, P. (1995): Susceptibility of safflower aphid, *Dactynotus carthami* HRL to some commonly used insecticides with a note on safety margin for the predator (*Coccinella septempunctata*) and honey bee (*Apis cerana*).

- Singh, D.S. and Yadav, T.D. (1994): Toxicity of deltamethrin, chloryrifosmethyl, etrimfos, malathion and fluvalinate against *Sitophilus oryzae* Linn. and *Trogoderma granarium* Everts. Indian J. Ent., **56**(4): 322-325.
- Singh, D.S.; Sircar, P.; Srivastava, V.S. and Dhingra, S. (1985): Biological efficacy of botanical products against some important insect pests. Indian J. Ent., **47**(4): 444-451.
- Singh, G. and Misra, P.N. (1988): Efficacy of new insecticides for control of insect pests of spring okra (*Abeinzoschus esculentus*). Ind. J. Agri. Sci., **58**(10): 783-785.
- Singh, G. and Singh, H. (1975): Biological studies on *Heliothis armigera* (Hubner) in the Punjab. Indian J. Ent. **37**(2): 154-164.
- Singh, G.P. and Goel, S.C. (1987): Body chaetotaxy of three noctuid caterpillars (Lepidoptera). Disch. Ent. Z., N.F. 34, (4-5): 373-381.

- Singh, H.; Katra, V.K. and Rohilla, H.R. (1986): Studies on the efficacy and economics of sprays of synthetic pyrethroids against mustard aphid, *Lipaphis erysimi* (Kalt.) on rapeseed. Ind. J. Plt. Prot., **14**(1): 1-5.
- Singh, H.; Mrig, K.K. and Mahler, J.C. (1996): Efficacy and persistence of products against lesser grain borer. *Rhyzopertha dominica* (F.) in wheat grains. Annals Biology Ludhiana, **12**(1): 99-103.
- Singh, H.M.; Singh, R. and Rizvi, S.M.A. (1987): Screening of synthetic pyrethroids against *Heliothis armigera* attacking chickpea. J. Agri. Res., **2**(2): 140-143.
- Singh, O.P. and Rawat, R.R. (1983): Seasonal incidence and toxicological studies in *Lipaphis erysimi* (Kalt.) and its parasite *Aphididus* sp. in Madhya Pradesh India. In the Aphids ed. B.K. Bahura. Zool. Soc. of Orissa. Utkal Univ. Bhubaneshwar, India, p.259-267.

- Singh, R.P. (1993): In Botanical Pesticides in developing countries; current status and future trends (eds. Dhaliwal, G.S. and Singh, B.), Common Wealth Publishers, New Delhi, p.236-268.
- Singh, S.P.; Raghavendra, K.; Singh, R.K. and Subbarao, S.K. (2001): Studies on larvicidal properties of leaf extract of *Solanum nigrum* L. (family Solanaceae). Current Science, **81**(12): 1529-1530.
- Singh, Y. and Bichoo, S.L. (1976): Bionomics of spodoptera Exigua (Hb.) on gram. Indian J. Ent. **38**(2): 138-141.
- Singhal, R.N., Vats, L.K. and Singh, J.S. (1976): Food energy budget for the AK grasshopper, *Poecilocerus pictus* Fabr. (Acrididae: Orthoptera). Indian J. Ecol. **3**(2): 119-124.
- Sinha U.S.P., Bajapai, C.M., Brahmachari, B.N., Sinha, B.R.R.P. and Sinha, S.S. (1998): "Consumption and utilization of *Tarminalia arjuna* leaves in different instars of tropical

tasar silkworm *Autheracea mylitta*, Drury
sericologia, **38**: 483.

Sinha, S.N.; Chakrabarti, A.K.; Agnihotri, N.P.; Jain,
H.K. and Gajbhiye, V.T. (1983): Field
evaluation of synthetic pyrethroids against
thrip, *Thrips tabaci* (Lind.) on onion and their
residues. Indian J. Ent., **45**(4): 420-423.

Smith J.M. and Hoy, M.A. (1995): Rearing methods for
ageniaspis citricola(Hymenoptera: Encyridae)
and *Cirrospilus quadristratus* (Hymenoptera:
Eulophidae) released in a classical biological
control programme for the citrus leaf miner
Phyllocnistis citrella (Lepidoptera:
Gracillariidae). Florida Entomologist **78**: 600-
608.

Smith, J.M. and Hoy, M.A. (1995): Rearing techniques for
Ageniaspis citrichola and *phyllocnistis citrella*
in: M.A. Hoy, Ed. managing the citrus leaf
miner proceedings from an international
conference, Orland Florida, April 23-25, 1996,

p. 99.

Smith, J.T.; Ingles, J.P. and Soderlund, M.D. (1988):
Actions of the pyrethroid insecticides
cismethrin and cypermethrin on the
pyrethroid insecticides cismethrin expressed
in xenopus oocytes. Archives of Insect
Biochemistry and Physiology, **38**(3): 126-136.

Snodgrass, R.E. (1928): Morphology and evolution of the
insect head and its appendages. Smithsonian
Misc. Coil. **81**(3): 1-158.

Sohail Ahmed and Naima Din. (2009): Leaf powders of
Baril (*Ocimum basilicum* L.) Lantana (*Lantana
camara* L.) and Gardenia (*Gardenia
jasminoedes* Ellis) affect biology of
Callosobruchus chinensis L. (Coleoptera:
Bruchidae) Pak. Entomol., **31**.

Solomon, J.D. (1962): Characters for determining sex in
elm spanworm pupae. J. Econ. Ent., 55: 269-
270.

- Solunke, B.R. and Deshpande, S.V. (1991): Studies on use of plants products for the control of lemon butterfly larvae. J. Maharashtra Agric. Univ., **16**(2): 302-303.
- Songkittisunton, V. (1989): The efficacy of neem oil and neem extracted substances on the rice. J. National Research Council of Thailand, **21**(2): 37-58.
- Soo Hoo, C.F. and Fraenkel, (1966): The consumption, digestion and utilization of food plants by polyphagous insect, *Prodenia eridania* (Gramer) J. Insect Physiol **12**: 711-730.
- Sparks, T.C.; Shoui, H.H. and Wellenmeyer, E.G. (1982): Temperature toxicity relationships of pyrethroids on three lepidoptera. J. Econ. Entomol., **75**: 643-644.
- Sreedharan, K.; Ghacko, M.J.; Rao, L.V.A. and Rao, L.G. (1981): Compatibility and bioefficacy of three synthetic pyrethroid insecticides with Bordeaux mixture. J. of Coffee Research,

11(3): 100-104.

Srivastava, B.K. and Mathur, L.M.L. (1964): Morphology and musculature of the head capsule of mature larva of *Prodenia litura* Fabr. (Lepidoptera: Noctuidae) Indian J. Ent. **26**(1): 78-91.

Srivastava, K.M.; Srivastava, S.K. and Singh, D.R. (1991): Efficacy of selected insecticides against mustard aphid, *Lipaphis erysimi* Kalt. (Homoptera: Aphididae). Entromon, **16**: 241-244.

Srivastava, R.P. and Pandey, S.N. (1967): On the immature stages of *Tarucus theophrastus* F. (Lepidoptera: Lycaenidae) a pest of ber (*Zizyphus* sp.). Proc. Nat. Acad. Science, **37**(B): 209-213.

Stein, U. and Klingauf, F. (1990): Insecticidal effect of plant extracts from tropical and subtropical species. Traditional methods are good as long as they are effective. J. Appl. Ento., **110**(2):

160-166.

Stickney, F.S. (1923): The head capsule of Coleoptera.

Illinois Biol. Mongr. **8**: 1-105.

Subramanian, S. and Nachiappan, R. (1996): Insect growth regulatory activity of aqueous extracts of *Ocimum basilicum* on rice brown plant hopper *Nilaparvata lugens*. Madras Agric. J., **83**(3): 200-201.

Sudhakar, K. (1991): In effectiveness of insecticides to control *Helicoverpa armigera* (Hubner) larva on cotton. Ind. J. Plt. Prot., **19**(2): 213-214.

Sukul, P.; Handa, S.K. and Singh, K.M. (1987): Comparative efficacy of synthetic pyrethroids against pod borer complex, crop growth and yield of green gram. Indian J. Ent. **49**(3): 408-413.

Sukumar, Kumuda, Perich, Michael, J. and Boobar, Lewis R. (1991): J. Am. Mosq. Control Assoc., **7**: 210-237.

Sundari, M.S.N.; Chockalingam, S. and Durairaj, G. (1999): Synergistic activities of plant extracts on a polyphagous lepidopteran pest, *Euproctis fraternus*. Journal of Environmental Biology, **20**(3): 245-257.

Syed Shayfur Rahman, Md. Mizanur Rahman, Md. Mizanur Rahman, Shameem Ara Begum, Balaram Roy and S.M. Fakruddin Shahed (2007): Ethanolic extract of melgota (*Macaranga postulate*) for repellency, insecticidal activity against rice weevil. African J. of Biotechnology, **6**(4): 379-383.

Syed Shayfur Rahman, Md. Mizanur Rahman, Shameem Ara Begum, Md. Mizanur Khan and Md. Mosharef Hossain Bhuiyan (2007): Investigation of *Sapindus mukorossi* extracts for repellency, insecticidal activity and plant growth regulatory effect. J. App. Sci. Res., **3**(2): 95-101.

Tandi, B.L.; Yadava, C.P.S. and Bhatnagar, K.N. (1993):

Bioefficacy of various insecticides against Red Hairy caterpillar, *Amsacta Moorei* Butler. Indian J. Ent., **55**(3): 317-324.

Tanzubil, P.B., G.W.K. Mansah, *et al.* (2000): Diapaus initiation and incidence in the millet stem borer, *Coniesta ignefusalis* (Lepidoptera: Pyralidae). The role of the host plant. Bulletin of entomological research (Print) August **90**(4): 365-371.

Thakar, A.V. and Srivastava, R.P. (1983): Chaetotaxy of eight noctuid caterpillars. Bull. Ent. **24**(2): 83-94.

Thakar, A.V.; Ameta, O.P.; Tripathi, N.N. and Vyas, A. (1992): Relative efficacy of indigenous plant extract gram pod borer *Heliothis armigera* (Hub.). Proceedings, first National Symposium, Allelopathy in agro-ecosystems (Agriculture and Forestry). Feb. 12-14, held at C.C.S. Haryana Agricultural Univ. Hisar, India (Edt. by Toro, P., Narwal, S.S. 162, Hisar India).

Ind. Soc. Allelopathy.

- Thakur, R.C.; Nema, K.K.; Kango, K.N. (1988): Comparative efficacy of neem seed kernel and some insecticidal formulations against the gram pod borer, *Heliothis armigera* (Hubner). Legume Research, **11**(3): 114-116.
- Thobbi, V.V. (1961): Growth potential of *Prodenia litura* F. in relation to certain food plants at Surat, Indian J. Ent.**23**(4): 262-264.
- Thomas, J. and Phadke, K.G. (1992): Field efficacy of chloropyriphos and quinalphos EC and dusts as compared to oxydemeton-methyl EC against aphid. Indian J. Entomol., **54**(2): 150-
- Thomas, J. and Phadke, K.G. (1993): Residual toxicity and persistence of chloropyriphos, quinalphos, and oxydemeton-methyl to aphid, *Lipaphis erysimi* (Kalt.). Indian J. Entomol., **55**: 275-280.
- Thomas, T.G.; Sharma, S.K.; Jalees, S. and Rahman,

- S.J. (1994): J. Basic Appl. Biomed., **2**: 53-55.
- Tiwari, R. and Dixit, V. (1994): Fungitoxic activity of vapour of some higher plants against predominant storage fungi. National Academy Science Letters, **17**(3-4): 55-57.
- Tompkins, G.J. Linduska, J.J., Young J.M. and Dougherty E.M. (1986): Effectiveness of microbial and chemical insecticides for controlling cabbage looper (Lepidoptera: Noctuidae) and imported cabbage worm (*Lepidoptera pieridae*) on collards in Maryland Journal of Economic Entomology, **79**: 497-501.
- Tripathi , A.K.; Srikant Sharma, Sushil Kumar and Edison, S. (ed.), Ramana, K.V. (ed.); Sasikumar, B. (ed.); Babu, K.N. (ed.) and Eapen, S.J. (1997): Repellent and insecticidal properties of piper retrofractum against insect pest of crops and stored grain. Biotechnology of species, medicinal and aromatic plants.

Proceedings. In National Seminar on
Biotechnology of Species and Aromatic Plants,
Calicut, India, 24-25 April, 1996, p.134-138.

Tripathi, Y.C. (2000): Potential of neem constituents in
pest management. *Everyman's Science*. **35**(2):
78-84.

Trivedi, P.C. (2002): *Plant Pest Management*. Aaviskar
Publishers, Jaipur.

Tsou, T.H. (1914): Homology of the body setae of some
lepidopterous larvae. *trans. Amer. micro,
Soc.* **33**: 223-260.

Vallella P. and Scriver J.M. (2002): Latitudinal variation
in photoperiodic induction of pupal diapause
in the spice bush swallowtail butterfly, *Papilio
trailus* Holarctic Lepidoptera **8**: 37-41.

Vas. R. de; de. Vas. R. (2003): Migrating lepidoptera in
2000 and recent adventive records. Sixty first
report. *Entomologische. Berichten*. **63**: 1, 14-
20; 6.

- Vats, L.K. and Kaushal, B.R. (1981): Population dynamics, recondary productivity and energy budget of *Parahieroglyphus bilineatus* Bol. (Orthoptera: Acrididae: Catantopinae). Acta Oecologia OECOL Gener.**2**: 355-369.
- Vats, L.K., Singh, J.S. and Yadava, P.S. (1977): Food energy budget of *Pieris brassicae* larvae - A pest of cruciferous agro-Ecosystems. Agro-Ecosystems, **3**: 303- 312.
- Vekaria, M.V. and Patel, G.M. (2000): Bioefficacy of Botanicals and certain chemical insecticides and their combinations against the mustard aphid, *Lipaphis erysimi*. Indian J. Ent., **62**(2): 150-158.
- Verma, R.K., Vinod K. Rajput and N. Ullah (2005): Light trapping of *Helicoverpa armigera* in the control of migratory pest of agriculture. VII National symposium on Advances in Indian Entomology: Productivity & Health organized by UttarPradesh Zoological Society2-4

October.

Verma, S. (1980): Evaluation of insecticides against pests and predators of the mustard crop. Indian J. Entomol., **42**: 582-591.

Vimala, R.; Shivaprakasam, K. and Seetharaman, K. (1993): Effect of plant extracts on the incidence of brinjal wilt. Madras Agric.J., **80**(7):409-412.

Waldbauer, G.P. (1964): The consumption, digestion and utilization of solanaceous and non-solanaceous plants by larvae of tobacco horn worm, *Protoparce sexta* (Johan) (Lepidoptera: Sphingidae). Entomol. Exp. Appl. **7**: 253-269.

Waldbauer, G.P. (1968): The consumption and utilization of food by insects. Adv. Insect. Physiol. **5**: 229-288. Acad. Press, N.Y. (Beament, Trcherne and Wigglesworth, Eds.).

Ward, K.E. and Landolt, P.J. (1955): Influence of Multiple Matings on fecundity & Longevity of

Female cabbage looper moths (Lepidoptera: Noctuidae) Ann. Entomol. Soc. Ann. **88**(6): 768-772.

White, N.D.G. and G.J. Demianyk (1999): the Bioenergetics of insects feeding on stored cereals. PP. 1-20 in C.A. Barlow (Ed.), Insects population growth and the environment. Research Sighpost, Trivendrum, India.

Williams, Carrol, M. (1970): Hormonal interactions between plants and insects. In: Ernest Sondheimer and John B. Simeone. Eds. Chemical Ecology, New York, London, Academic Press, pp.103-129.

Yadav, P.R., Yadav, C.R. and Sachan J.N. (1983): Influence of temperature on the development of cabbage semi looper, *Trichoplusia ni* Hub Bulletin of Entomology. **24**: 54-57.

Yadav, R.L. (1972): Life history and economic importance of *Polytela gloriosae* Fabr. (Hadeninae: Noctuoidae), Lepidoptera. Bull. Ent. **13**(1): 14-

18.

Yadav, T.D. (1987): Toxicity of deltamethrin, cypermethrin, permethrin against thirteen stored product insects. Indian J. Ent., **49**(1): 21-26.

Youm, O. and E.O. Owusu (1998): Assessment of yield loss due to the millet head miner, *Heliochcilus albipunctella* (Lepidoptera: Noctuidae) using a damage rating scale and regression analysis in Niger. International journal of Pest management **44**(2): 119-121.

Zalucki M.P., Clarke A.R. and Malcolm S.B. (2002): Ecology and behaviour of first instar larval Lepidoptera. Annu. Rev. Entomol. **47**: 361-393.