

## LIST OF TABLES

<i>Table No.</i>	<i>Titles</i>	<i>Page No.</i>
<b>Table No. 1.1</b>	<b>Comparison between India and World Maternal and Infant Mortality rate</b>	<b>5</b>
<b>Table No. 3.1</b>	<b>Lucknow District and Lucknow Urban in Census 2011</b>	<b>54</b>
<b>Table No. 3.2</b>	<b>Demographic profile of Lucknow City</b>	<b>54</b>
<b>Table No. 3.3</b>	<b>Health Facility in Mawaiya slum</b>	<b>55</b>
<b>Table No. 4.1</b>	<b>Age of the respondents</b>	<b>156</b>
<b>Table No. 4.2</b>	<b>Literacy status</b>	<b>156</b>
<b>Table No. 4.3</b>	<b>School going Status</b>	<b>156</b>
<b>Table No. 4.4</b>	<b>Educational Status</b>	<b>156</b>
<b>Table No. 4.5</b>	<b>Type of family</b>	<b>157</b>
<b>Table No. 4.6</b>	<b>Religion</b>	<b>157</b>
<b>Table No. 4.7</b>	<b>Caste/ Sub-caste</b>	<b>157</b>
<b>Table No. 4.8</b>	<b>Monthly income of the Family</b>	<b>157</b>
<b>Table No. 4.9</b>	<b>Type of House</b>	<b>158</b>
<b>Table No. 4.10</b>	<b>Source of Drinking Water at Home</b>	<b>158</b>
<b>Table No. 4.11</b>	<b>Latrine facility in House</b>	<b>158</b>
<b>Table No. 4.12</b>	<b>Occupation</b>	<b>158</b>
<b>Table No. 4.13</b>	<b>Source for Cooking Food</b>	<b>159</b>
<b>Table No. 4.14</b>	<b>Age at Marriage</b>	<b>159</b>
<b>Table No. 4.15</b>	<b>Pregnancy registered with the ANM, AWW, ASHA or other health provider</b>	<b>159</b>
<b>Table No. 4.16</b>	<b>Pregnancy registration</b>	<b>160</b>
<b>Table No. 4.17</b>	<b>Month of pregnancy registration</b>	<b>160</b>
<b>Table No. 4.18</b>	<b>Registration card obtained</b>	<b>160</b>
<b>Table No. 4.19</b>	<b>Receive antenatal care/check up during the pregnancy</b>	<b>161</b>
<b>Table No. 4.20</b>	<b>Place of antenatal care</b>	<b>161</b>
<b>Table No. 4.21</b>	<b>Number of times antenatal check up was received</b>	<b>162</b>
<b>Table No. 4.22</b>	<b>Quantity of food intake increased during pregnancy</b>	<b>162</b>

<b>Table No. 4.23</b>	<b>Period of increased food intake</b>	<b>162</b>
<b>Table No. 4.24</b>	<b>Preparation for Delivery</b>	<b>163</b>
<b>Table No. 4.25</b>	<b>TT injection administered</b>	<b>163</b>
<b>Table No. 4.26</b>	<b>Month of pregnancy when TT vaccination was administered</b>	<b>164</b>
<b>Table No. 4.27</b>	<b>Receive Iron Folic Acid tablets or syrup</b>	<b>164</b>
<b>Table No. 4.28</b>	<b>Supplementary nutrition received from Anganwadi center during this pregnancy</b>	<b>164</b>
<b>Table No.4.29</b>	<b>Supplementary nutrition from Anganwadi center was regularly received</b>	<b>164</b>
<b>Table No. 4.30</b>	<b>Consumption of the nutrition received from the Anganwadi center</b>	<b>165</b>
<b>Table No. 4.31</b>	<b>Regular afternoon rest during the last three months of pregnancy</b>	<b>165</b>
<b>Table No. 4.32</b>	<b>Place of Delivery</b>	<b>165</b>
<b>Table No. 4.33</b>	<b>Assistance at the time of delivery</b>	<b>166</b>
<b>Table No. 4.34</b>	<b>Instrument used to cut the Umbilical Cord Percent</b>	<b>166</b>
<b>Table No. 4.35</b>	<b>Tie the cord</b>	<b>166</b>
<b>Table No. 4.36</b>	<b>Surface used to cut the Umbilical cord</b>	<b>167</b>
<b>Table No. 4.37</b>	<b>Ungent applied on the cord</b>	<b>167</b>
<b>Table No. 4.38</b>	<b>Factors for choosing Hospital for Delivery</b>	<b>167</b>
<b>Table No. 4.39</b>	<b>Mode of transport was used to go to the hospital</b>	<b>168</b>
<b>Table No. 4.40</b>	<b>Escort to the Hospital</b>	<b>168</b>
<b>Table No. 4.41</b>	<b>Payment of fee for services during delivery</b>	<b>168</b>
<b>Table No. 4.42</b>	<b>Personnel who did the delivery</b>	<b>168</b>
<b>Table No. 4.43</b>	<b>Type of delivery in hospital</b>	<b>169</b>
<b>Table No. 4.44</b>	<b>Number of days stay in the hospital</b>	<b>169</b>
<b>Table No. 4.45</b>	<b>Time taken to clothe the Neonatal</b>	<b>169</b>
<b>Table No. 4.46</b>	<b>Bathe the baby on the day of birth</b>	<b>170</b>
<b>Table No. 4.47</b>	<b>Bathing Frequency</b>	<b>170</b>
<b>Table No. 4.48</b>	<b>Methods to keep the baby warm at home</b>	<b>170</b>
<b>Table No. 4.49</b>	<b>Baby cry immediately after birth</b>	<b>170</b>

<b>Table No. 4.50</b>	<b>Religious practices followed after child birth</b>	<b>171</b>
<b>Table No. 4.51</b>	<b>Methods to clear the Airway</b>	<b>171</b>
<b>Table No. 4.52</b>	<b>Breastfeeding begin time</b>	<b>171</b>
<b>Table No. 4.53</b>	<b>Reasons for late initiation of breast feeding</b>	<b>171</b>
<b>Table No. 4.54</b>	<b>Colostrum given to the baby</b>	<b>172</b>
<b>Table No. 4.55</b>	<b>Bottle feed the baby</b>	<b>172</b>
<b>Table No. 4.56</b>	<b>Reasons for bottle feeding the baby</b>	<b>172</b>
<b>Table No. 4.57</b>	<b>Nutriments besides breast milk giving to the infant</b>	<b>173</b>
<b>Table No. 4.58</b>	<b>Breast feed when you are ill</b>	<b>173</b>
<b>Table No. 4.59</b>	<b>Polio vaccine at the time of birth</b>	<b>173</b>
<b>Table No. 4.60</b>	<b>BCG vaccine given to the neonate</b>	<b>173</b>
<b>Table No. 4.61</b>	<b>The child was treated when unwell</b>	<b>174</b>
<b>Table No. 4.62</b>	<b>Home remedies given to the neonate in case of stomach ache</b>	<b>174</b>
<b>Table No. 4.63</b>	<b>Home remedies to treat diarrhea</b>	<b>174</b>
<b>Table No. 4.64</b>	<b>Home remedies to treat cold and cough</b>	<b>175</b>
<b>Table No. 4.65</b>	<b>Infant given massages</b>	<b>175</b>
<b>Table No. 4.66</b>	<b>Lubricants used to massage the infant</b>	<b>175</b>
<b>Table No. 4.67</b>	<b>Frequency of massage</b>	<b>175</b>
<b>Table No. 4.68</b>	<b>Methods to clean the airway for normal breathing of the neonate</b>	<b>176</b>
<b>Table No. 4.69</b>	<b>Started Breastfeeding</b>	<b>176</b>
<b>Table No. 4.70</b>	<b>Advantages of giving colostrums to the baby</b>	<b>176</b>
<b>Table No. 4.71</b>	<b>No. of months baby should be breast fed only</b>	<b>176</b>
<b>Table No. 4.72</b>	<b>Reasons for feeding the infant in illness</b>	<b>177</b>
<b>Table No. 4.73</b>	<b>Signs when the baby is seriously ill</b>	<b>177</b>
<b>Table No. 4.74</b>	<b>Danger Sign</b>	<b>177</b>
<b>Table No. 4.75</b>	<b>Advantages of massage</b>	<b>178</b>
<b>Table No. 4.76</b>	<b>Mothers know by themselves how to breast feed their child, so counseling is not required</b>	<b>178</b>
<b>Table No. 4.77</b>	<b>For the first 6 months of life baby should not be given water, “JanamGhuti,” honey or any other milk except</b>	

	<b>breast milk</b>	<b>178</b>
<b>Table No. 4.78</b>	<b>Immunizing the child is important as this will save the child from many diseases</b>	<b>179</b>
<b>Table No. 4.79</b>	<b>Breast milk provides nourishment to the newborn</b>	<b>179</b>
<b>Table No. 4.80</b>	<b>Wash hands thoroughly with soap and water before handing the newborn</b>	<b>179</b>
<b>Table No. 4.81</b>	<b>Breast feeding helps in forming an emotional bond between the mother and the newborn</b>	<b>180</b>
<b>Table No. 4.82</b>	<b>Massaging with oil before bath helps to keep the newborn skin healthy and improves blood circulation</b>	<b>180</b>
<b>Table No. 4.83</b>	<b>Child should always be taken to a big hospital if he/she falls sick</b>	<b>180</b>
<b>Table No. 4.84</b>	<b>The newborn should be immunized during the first week of life with BCG,OPV, and hepatitis B</b>	<b>181</b>
<b>Table No. 4.85</b>	<b>Children get loose motions due to effect of evil eye</b>	<b>181</b>
<b>Table No. 4.86</b>	<b>Colostrum is harmful to the newborn</b>	<b>181</b>
<b>Table No. 4.87</b>	<b>The newborn should be bathed immediately after birth</b>	<b>181</b>
<b>Table No. 4.88</b>	<b>The practice of applying kajal to the newborn eyes is good</b>	<b>182</b>
<b>Table No. 4.89</b>	<b>Newborn should not be dressed with soft and cleaned clothes</b>	<b>182</b>
<b>Table No. 4.90</b>	<b>Breast milk is not best food for newborn</b>	<b>182</b>
<b>Table No. 4.91</b>	<b>A baby less than 2.5 kg will definitely gain weight and so needs no special care</b>	<b>182</b>