

LIST OF TABLES

TABLE NO.	TITLE	PAGE NO.
2. REVIEW OF LITERATURE		
2.1	Characteristic features of white rot fungi along with brown and soft rot wood decaying fungi	10
2.2	Exopolysaccharide producing white rot fungi versus other microorganisms	15
2.3	Industrial and medicinal application of microbial exopolysaccharides	17
2.4	Examples of homopolysaccharides along with their composition and functions	20
2.5	Examples of heteropolysaccharides along with their composition and functions	21
2.6	Summarizes microorganisms producing microbial flocculants	56
2.7	Flocculation mechanism for different types of flocculants	58
2.8	List of commercial thickening agent with their sources and applications.	63
2.9	Other important commercial applications of exopolysaccharides produced by different microbial types.	65
3. MATERIALS & METHODS		
3.1	General Information & Macroscopic details observed during collection	89
3.2	Parameters studied for microscopic features of the various collected white rot samples	90
3.3	Parameters studied for cultural features of the various collected white rot samples:	90
3.4	Plackett Burman design for screening variables at different level of ranges for EPS production	98
3.5	Experimental range and levels of independent factors for RSM design.	99
4. RESULTS & DISCUSSION		
4.1	List of white rot fungal species collected from various biodiverse regions of Chandigarh capital region	121
4.2	The generic characteristics of 16 different white rot fungi collected	124

	from biodiverse locations of Chandigarh and nearby regions.	
4.3	Taxonomic description of <i>Pleurotus</i> sp. (RDM9)	130
4.4	Aligned sequence data of <i>Pleurotus pulmonarius</i> RDM9	133
4.5	Closest species along with their GenBank accession number and % identity from BLASTn search	133
4.6	Parameters, experimental runs and responses (EPS production) of Plackett-Burman Design used for the screening of significant parameters for EPS production under submerged fermentation by <i>Pleurotus pulmonarius</i> RDM9	152
4.7	Analysis of variance (ANOVA) for Plackett-Burman Design	153
4.8	Central Composite Design (CCD) matrix with experimental and predicted values of EPS production by <i>Pleurotus pulmonarius</i> RDM9	154
4.9	Analysis of variance (ANOVA) for Response Surface Methodology developed for optimum EPS yield by <i>Pleurotus pulmonarius</i> RDM9	156
4.10	Validation of EPS production in submerged fermentation with the predicted values	164
4.11	Final optimized conditions for EPS production	165
4.12	Solubility of the test EPS in various organic and inorganic solvents	179
4.13	The various parameters studied for organoleptic properties of EPS and standard pullulan	181
4.14	Summary of the functional groups present in the test EPS	183
4.15	Enlisting the peaks of XRD spectrum of the EPS	191
4.16	Mean ALT and AST of the control and EPS treated group	194
4.17	Emulsification activity (EA%) after 1 hour and emulsification index after 24, 48, 72 and 96 hours (%E ₂₄ , %E ₄₈ , %E ₇₂ and %E ₉₆) of the test EPS and the standard Guar gum with different vegetable oils and hydrocarbons	197
4.18	The effect of different dosage of the test EPS on the emulsification activity (%EA)	198
4.19	Densities of emulsions formed by different EPS concentrations	200
4.20	State of the emulsion containing different concentrations of the test EPS	202
