

## REFERENCES

- Abu-Abed, S., Doll'e, P., Metzger, D., Beckett, B., Chambon, P. and Petkovich, M. (2001).** The retinoic acid metabolizing enzyme, CYP26A1, is essential for normal hindbrain patterning, vertebral identity, and development of posterior structures. *Genes Development* **15**:226–40.
- Abu-Abed, S., MacLean, G., Fraulob, V., Chambon, P., Petkovich, M. and Doll'e, P. (2002).** Differential expression of the retinoic acid-metabolizing enzymes CYP26A1 and CYP26B1 during murine organogenesis. *Mechanism of Development* **110**:173–177.
- Altucci, L., Leibowitz, M.D. and Ogilvie, K.M. (2007).** RAR and RXR modulation in cancer and metabolic disease. *Nature Review Drug Discovery* **6**:793–810.
- Altschul S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. (1990).** Basic Local Alignment Search Tool, *Journal of Molecular Biology* **215**: 403-410.
- Bajorath, J. (2002).** Integration of virtual and high-throughput screening. *Nature Review Drug Discovery* **1**:882–894.
- Baudet, A., Quere, R., Cassinat, B., Bertrand, G., Marti, J., Manchon, L., Piquemal, D., Chomienne, C. and Commes, T. (2007).** Pharmacogenomic analysis of acute promyelocytic leukemia cells highlights CYP26 cytochrome metabolism in differential all-trans retinoic acid sensitivity. *Blood*. **109**: 4450-4460.
- Berendsen, H.J.C., Postma, J.P.M., Gunsteren, W.F., DiNola, A. and Haak, J.R. (1984).** Molecular dynamics with coupling to an external bath. *The Journal of Chemical Physics* **8**: 3384-3690.
- Blomhoff, R. and Blomhoff, H.K.(2006).** Overview of retinoid metabolism and function. *Journal of Neurobiology* **66**:606–630.

**Bowles, J., Knight, D., Smith, C., Wilhelm, D. and Richman, J. (2006).** Retinoid signaling determines germ cell fate in mice. *Science* **312**:596–600.

**Bowie, J. U., Luthy, R. and Eisenberg, D. (1991).** A method to identify protein sequences that fold into a known three-dimensional structure. *Science* **253**:164–170.

**Blundell, T.L., Sibanda, B.L., Sternberg, M.J.E. and Thornton, J.M. (1987).** Knowledge-based prediction of protein structures and the design of novel molecules. *Nature* **326**:347–352.

**Catharine, R., Ross, A. and Zolfaghari, R. (2004).** Regulation of Hepatic Retinol Metabolism: Perspectives from Studies on Vitamin A Status 1,2. *Journal of Nutrition* **134**: 269S–275S.

**Catharine, R., Ross, A. and Reza, Z. (2011).** Cytochrome P450s in the Regulation of Cellular Retinoic Acid Metabolism. *Annual Review. Journal of Nutrition.* **31**:65–87.

**Chambon P. (1996).** A decade of molecular biology of retinoic acid receptors. *FASEB Journal.* **10**:940–954.

**Chapman, J. S., Weiss, K. L., Curley, R.W.J., Highland, M.A. and Clagett-Dame, M. (2003).** Hydrolysis of 4-HPR to atRA occurs in vivo but is not required for retinamide-induced apoptosis. *Archives of Biochemistry and Biophysics.* **419**:234–243.

**Chen, Z. and DeTh'e, H. (2010).** Acute promyelocytic leukaemia: novel insights into the mechanisms of cure. *Nature Review:Cancer* **10**:775–783.

**Chris, O., Alessandra, V., Alan, E. and Wilfred, F.V.G. (2004).** A biomolecular force field based on the free enthalpy of hydration and solvation the GROMOS force-field parameter sets 53A5 and 53A6. *Journal of Computational Chemistry* **13**: 1656–1676.

**Clagett-Dame, M. and DeLuca, H.F. (2002).** The role of vitamin A in mammalian reproduction and embryonic development. *Annual Review of Nutrition*. **22**:347–381.

**Colovos, C. and Yeates, T.O. (1993).** Verification of protein structures: Patterns of nonbonded atomic interactions. *Protein Science* **2**:1511–1519.

**Croce, C.M. (2008).** Oncogenes and Cancer. *The New England Journal of Medicine* **358** : 502-511.

**Danielson, P.B. (2002).** The cytochrome P450 superfamily: biochemistry, evolution and drug metabolism in humans. *Current Drug Metabolism* **3**:561-597.

**David, W. and Rithcie, D.W. (2000).** Evaluation of Protein Docking Predictions using Hex 3.1 in CAPRI round 1-2. *Protein, Structure, Function and Genetics*, Wiley-liss Inc.

**DeLuca, H.F. (1979).** Retinoic acid metabolism . *Federation Proceeding* **38**:2519–2523.

**DeLuca, H.F. and Roberts, A.B. (1969).** Pathways of retinoic acid and retinol metabolism. *American Journal of Clinical Nutrition* **22**:945–952.

**De Marco, P., Elisa Merello, E., Mascelli, S., Raso, S., Santamaria, A., Ottaviano, C., Calevo, M. G., Cama, A. and Capra, V. (2006).** Mutational Screening of the *CYP26A1* Gene in Patients with Caudal Regression Syndrome. *Wiley-Liss, Inc. Birth Defects Research*. **76**:86–95.

**Duester, G. (2008).** Retinoic acid synthesis and signaling during early organogenesis. *Cell* **134**:921–931.

**Eckhoff, C., Collins, M.D. and Nau, H. (1991).** Human plasma all-trans-13-cis- and 13-cis-4-oxoretinoic acid profiles during subchronic vitamin A supplementation: comparison to retinol and retinyl ester plasma levels. *Journal of Nutrition* **121**:1016–1025.

**Eckhoff, C. and Nau, H. (1990).** Identification and quantitation of all-trans- and 13-cis-retinoic acid and 13-cis-4-oxoretinoic acid in human plasma. *Journal of Lipid Research* **31**:1445–1454.

**Edwin, S., Christina, E., van den Brink, L., Tertoolen, G.J., Bart van der, B. and Paul, T. (1999).** Retinoic Acid Hydroxylase (CYP26) Is a Key Enzyme in Neuronal Differentiation of Embryonal Carcinoma Cells. *Developmental Biology* **213**: 390–404.

**Eisenberg, D., Luthy, R. and Bowie, J.U. (1997).** VERIFY3D: assessment of protein models with three-dimensional profiles. *Methods in Enzymology* **277**: 396–404.

**Essmann, U., Perera, L., Berkowitz, M.L., Darden, T., Lee, H. and Pedersen, L.G. (1995).** A smooth particle meshewald method. *Journal of Chemical Physics*. **103**: 8577–8593.

**Evan, G.I. and Vousden, K.H. (2001).** Proliferation, cell cycle and apoptosis in cancer. *Nature (Lond)* **411**:342–348.

**Fex, G. and Felding, P. (1984).** Factors affecting the concentration of free holo retinol-binding protein in human plasma. *European Journal of Clinical Investigation* **14**:146–9.

**Fische, E. (1894).** Einfluss der Configuration auf die wirkung der Enzyme. *Berichte der Deutschen Chemischen Gesellschaft* **27**:2985–2993.

**Fisher, G. J. and Voorhees, J. J. (1996).** Molecular mechanisms of retinoid actions in skin. *FASEB Journal* **10**:1002–1013.

**Frolik, C.A., Roberts, A.B., Tavela, T.E., Roller, P.P., Newton, D. L. and Sporn, M.B. (1979).** Isolation and identification of 4-hydroxy- and 4-oxoretinoic acid. *In vitro* metabolites of all-trans-retinoic acid in hamster trachea and liver. *Biochemistry* **18**:2092–2097.

**Fujii, H., Sato, T. and Kaneko, S. (1997).** Metabolic inactivation of retinoic acid by a novel P450 differentially expressed in developing mouse embryos. *EMBO journal* **16**:4163–4173.

**Gaemers, I. C., van Pelt, A. M., van der Saag, P. T. and Rooij, D. G. (1996).** All-trans-4-oxo-retinoic acid: a potent inducer of in vivo proliferation of growth-arrested A spermatogonia in the vitamin A-deficient mouse testis. *Endocrinology* **137**:479–485.

**Garnier, J., Gibrat, J. F. and Robson, B. (1996).** GOR secondary structure prediction method version IV. *Methods in Enzymology* **266**: 540-553.

**Gasteiger, E., Gattiker, A., Hoogland, C., Ivanyi, I., Appel, R.D. and Bairoch, A. (2003).** ExpPASy: The proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Research*. **13**: 3784–3788.

**Giltaire, S., Herphelin, F., Frankart, A., H´erin, M., Stoppie, P. and Poumay, Y. (2009).** The CYP26 inhibitor R115866 potentiates the effects of all-trans retinoic acid on cultured human epidermal keratinocytes. *British Journal of Dermatology*. **160**:505–513.

**Gozani, O., Boyce, M., Yoo, L., Karuman, P. and Yuan. J. (2002).** Life and death in paradise. *Nature: Cell Biology* **4**:159–162.

**Green, D. R. and Evan, G. I. (2002).** A matter of life and death. *Cancer Cell* **1**:19–30.

**Gill, S.C. and von Hippel, P.H. (1989).** Calculation of protein extinction coefficients from amino acid sequence data. *Annual Review of Biochemistry* **182**:319-326.

**Gomaa, M. S., Yee, S. W., Milbourne, C.E., Barbera, M. C., Simons, C. and Brancale, A. (2006).** Homology model of human retinoic acid metabolizing enzyme cytochrome P450 26A1 (CYP26A1): active site architecture and ligand binding. *Journal of Enzyme Inhibition Medicinal Chemistry* **21**:361–369.

**Geourjon, C. and Deleage, G. (1995).** SOPMA: Significant improvement in protein secondary structure prediction by prediction from alignments and joint prediction. *CABIOS*. **11**:681-684.

**Greer, J. (1981).** Comparative model-building of the mammalian serine proteases. *Journal of Molecular Biology* **153**:1027–1042.

**Gu, X., Xu, F. and Song, W. (2006).** A novel cytochrome P450, zebrafish Cyp26D1, is involved in metabolism of all-trans retinoic acid. *Molecular Endocrinology* **20**:1661–1672.

**Gu, X., Xu, F. and Wang, X. (2005).** Molecular cloning and expression of a novel CYP26 gene (cyp26d1) during zebrafish early development. *Gene Expression Patterns* **5**:733–739.

**Gudas, L.J., Sporn, M.B. and Roberts, A.B. (1994).** The Retinoids: Biology, Chemistry, and Medicine, (Sporn, M.B.; Roberts, A.B. and Goodman, D.S. Eds.). *Raven Press, New York*, pp. 443-520.

**Hanahan, D. and Weinberg, R. A. (2000).** The hallmarks of cancer. *Cell* **100**:57-70.

**Heeg, S., Doebele, M., Von Werder, A. and Opitz, O. G. (2006).** *In vitro* transformation models: modeling human cancer. *Cell Cycle* **5 (6)**: 630-634.

**Hess, B., Bekker, H., Berendsen, H.J.C. and Fraaije, J.G.E.M. (1997).** LINCS: A linear constraint solver for molecular simulations. *Journal of Computational Chemistry* **12**: 1463–1472.

**Hernandez, R.E., Putzke, A.P. and Myers, J.P. (2007).** CYP26 enzymes generate the retinoic acid response pattern necessary for hindbrain development. *Developmental Biology* **134**:177–187.

**Hess, B., Kutzner, C., Spoel, D. and Lindahl, E. (2008).** GROMACS 4: Algorithms for Highly Efficient, Load- Balanced, and Scalable Molecular Simulation. *Journal of Chemical Theory and Computation* **3**:435–447.

**Hong, E., Chang , C. I., Lao-Sirieix, P. and Fitzgerald, R.C. (2008).** A novel role for the retinoic acid-catabolizing enzyme CYP26A1 in Barrett's associated adenocarcinoma. *Oncogene*. **27**: 2951–2960.

**Ikai, A. J. (1980).** Thermostability and aliphatic index of globular proteins. *Journal of Biochemistry* **88**:1895-1898.

**Irwin, J. J. and Shoichet, B.K. (2005).** ZINC - A Free Database of Commercially Available Compounds for Virtual Screening. *Journal of Chemical Information and Modeling* **45** :177-182.

**Jacolot, F., Simon, I., Dreano, Y., Beaune, P., Riche, C. and Berthou, F. (1991).** Identification of the cytochrome P450 IIIA family as the enzymes involved in the N-demethylation of tamoxifen in human liver microsomes. *Biochemical Pharmacology*. **41**:1911–1919.

**Jayne, E., Thatcher and Isoherranen, N. (2009).** The role of CYP26 enzymes in retinoic acid clearance. *Expert Opinion Drug Metabolism Toxicology* **5**:875–886.

**Jayne, E. Thatcher., Brian Buttrick., Scott, A., Shaffer, Jakob A, Shimshoni., David R. Goodlett., Wendel L. Nelson. and Nina Isoherranen. (2011).** Cytochrome P450 mediated metabolic oxidation. *Molecular Pharmacology* **80**:228–239.

**John, K. V., Lakshmanan, M. R. and Cama, H. R. (1967).** Preparation, properties and metabolism of 5,6-monoepoxyretinoic acid. *Journal of Biochemistry* **103**:539–543.

**Jorgensen, W. L. (2004).** The many roles of computation in drug discovery. *Science* **303**: 1813–1818.

**Karlsson, M., Strid, A., Sirsjo, A. and Eriksson, L.A. (2008).** Homology models and molecular modeling of human retinoic acid metabolizing enzymes cytochrome P450 26A1 (CYP26A1) and P450 26B1 (CYP26B1). *Journal of Chemical Theory and Computational* **4**:1021–1027.

**Klaassen, I., Brakenhoff, R. H., Smeets, S. J., Snow, G. B. and Braakhuis, B. J. (2001).** Metabolism and growth inhibition of four retinoids in head and neck squamous normal and malignant cells. *British Journal of Cancer* **85**:630–635.

**Kelley, L.A. and Sternberg, M. J. (2009).** Protein structure prediction on the Web: a case study using the Phyre server. *Nature. Protocol* **4**:363–371.

**Krüger, K. A., Blum, J. W. and Greger, D. L. (2005).** Expression of nuclear receptor and target genes in liver and intestine of neonatal calves fed colostrum and vitamin A. *Journal of Dairy Science*. **88**:3971–3981.

**Kyte, J. and Doolittle, R. F. (1982).** A simple method for displaying the hydropathic character of a protein. *Journal of Molecular Biology*. **157**:105-132.

**Lane M.A., Chen A.C., Roman S.D., Derguini F. and Gudas, L.J. (1999).** Removal of LIF (leukemia inhibitory factor) results in increased vitamin A (retinol) metabolism to 4-oxoretinol in embryonic stem cells. *Proceeding of Natural Academy of Science USA*. **96**:13524–13529.

**Langton, S. and Gudas, L.J. (2008).** CYP26A1 knockout embryonic stem cells exhibit reduced differentiation and growth arrest in response to retinoic acid. *Developmental Biology*. **315**:331–354.

**Laskowski, R.A., MacArthur, M.W., Moss, D.S. and Thornton, J.M. (1993).** PROCHECK: a program to check the stereochemical quality of protein structures. *Journal of Applied Crystallography*. **26**: 283–291.



**Lee, G.H. and Osanai, M. (2011).** Enhanced expression of retinoic acid-metabolizing enzyme CYP26A1 in sunlight-damaged human skin. *Medical Molecular Morphology*. **44**:200-206.

**Linding, R., Russell, B. R., Neduva, V. and Gibson, J. T. (2003).** GlobPlot: exploring protein sequences for globularity and disorder. *Nucleic Acids Research*. **31**:3701-3708.

**Loudig, O., Babichuk, C., White, J., Abu-Abed, S., Mueller, C. and Petkovich, M. (2000).** Cytochrome P450RAI (CYP26) promoter: a distinct composite retinoic acid response element underlies the complex regulation of retinoic acid metabolism. *Molecular Endocrinology*. **14**:1483–1497.

**Loudig, O., Maclean, G.A., Dore, N.L., Luu, L. and Petkovich, M. (2005).** Transcriptional co-operativity between distant retinoic acid response elements in regulation of Cyp26A1 inducibility. *Journal of Biochemistry*. **392**:241–248.

**Lovell, S. S., Davis, I.W., Arendall III, W. B., Bakker, P.I.W., Word, J. M., Prisant, M. G., Richardson, J.S. and Richardson, D. C. (2002).** Structure validation by Ca geometry: F, C and Cb deviation. *Proteins: Structure Function & Genetics* **50**:437–450.

**Lutz, J. D., Dixit, V., Yeung, C. K., Dickmann, L. J. and Zelter, A. (2009).** Expression and functional characterization of cytochrome P450 26A1, a retinoic acid hydroxylase. *Biochemistry and Pharmacology*. **77**:258–268.

**MacLean, G., Abu-Abed, S., Dollé, P., Tahayato, A., Chambon, P. and Petkovich, M. (2001).** Cloning of a novel retinoic-acid metabolizing cytochrome P450, Cyp26B1, and comparative expression analysis with Cyp26A1 during early murine development. *Mechanism of Development*. **107**:195–201.

**MacLean G, Li H, Metzger D, Chambon P. and Petkovich, M. (2007).** Apoptotic extinction of germ cells in testes of Cyp26b1 knockout mice. *Endocrinology*. **148**:4560–4567.

**MacKerell, A.D. Jr., Bashford, D., Bellott, M., Dunbrack, R.L. Jr., Evanseck, J.D., Field, M.J., Fischer, S., Gao, J., Guo, H., Ha, S., Joseph-McCarthy, D., Kuchnir, L., Kuczera, K., Lau, F.T.K., Mattos, C., Michnick, S., Ngo, T., Nguyen, D.T., Prodhom, B., Reiher, W.E. III, Roux, B., Schlenkrich, M., Smith, J.C., Stote, R., Straub, J., Watanabe, M., Wiórkiewicz-Kuczera, J., Yin, D., and Karplus, M. (1998).** All-atom empirical potential for molecular modeling and dynamics studies of proteins. *Journal of Physical Chemistry B*. **102**:3586-3616.

**Makoto Osanai and Gang-Hong Lee (2011).** Enhanced expression of retinoic acid-metabolizing enzyme CYP26A1 in sunlight-damaged human skin. *Medical Molecular Morphology* **44**:200–206 .

**Mangelsdorf, D. J. and Evans, R.M. (1995).** The RXR heterodimers and orphan receptors. *Cell* **83**:841–850.

**Marill, J., Idres, N., Capron, C.C., et al (2003).** Retinoic acid metabolism and mechanism of action: A Review: *Current Drug Metabolism* **4**:1–10.

**McCaffery, P. and Simons, C. (2007).** Prospective teratology of retinoic acid metabolic blocking agents (RAMBAs) and loss of CYP26 activity. *Current Pharmaceutical Design*. **13**:3020–37

**McCormick, A. M., Napoli, J. L., Schnoes, H. K. and DeLuca, H. F. (1978).** Isolation and identification of 5, 6-epoxyretinoic acid: a biologically active metabolite of retinoic acid. *Biochemistry* **17**:4085–4090.

**McClure, M. A., Vasi, T. K. and Fitch, W. M. (1994).** Comparative analysis of multiple protein-sequence alignment methods. *Molecular Biology Evolution*. **11**:571-592.

**Mohamed, S. G., SookWah, Y., Elizabeth, M., Barbera, M.C., Claire, S. and Brancale, C. (2006).** Homology Model of Human Retinoic Acid Metabolising Enzyme Cytochrome P450 26A1 (CYP26A1): Active Site Architecture and Ligand Binding. *Journal of Enzyme Inhibition and Medicinal Chemistry* **21**: 361–369.

**Muindi, J., Frankel, S.R., Miller, W.H. Jr., Jakubowski, A., Scheinberg, D. A., Young, C. W., Dmitrovsky, E. and Warrell, R. P. Jr. (1992).** Continuous treatment with all-*trans* retinoic acid causes a progressive reduction in plasma drug concentrations: implications for relapse and retinoid “resistance” in patients with acute promyelocytic leukemia. *Blood* **79**:299–303.

**Nadin, L. and Murray, M. (1999).** Participation of CYP2C8 in retinoic acid 4-hydroxylation in human hepatic microsomes. *Biochemistry Pharmacology*. **58**:1201–1208.

**Napoli, J. L. (1999).** Retinoic acid: its biosynthesis and metabolism. *Program. Nucleic Acid Research : Molecular Biology*. **63**:139–188.

**Nau, H., Tzimas, G., Mondry, M., Plum, C. and Spohr, H. L.(1995).** Antiepileptic drugs alter endogenous retinoid concentrations: a possible mechanism of teratogenesis of anticonvulsant therapy. *Life Sciences*. **57**:53–60.

**Nelson DR. (1999).** A second CYP26 P450 in humans and zebrafish: CYP26B1. *Archives of Biochemistry and Biophysics*. **371**:345–347.

**Niederreither, K., Abu-Abed, S., Schuhbaur, B., Petkovich, M., Chambon, P. and Dollé, P. (2002).** Genetic evidence that oxidative derivatives of retinoic acid are not involved in retinoid signaling during mouse development. *Nature. Genetics*. **31**:84–88.

**Njar, V. C., Gediya, L., Purushottamachar, P., Chopra, P. and Vasaitis, T. S. (2006).** Retinoic acid metabolism blocking agents (RAMBAs) for treatment of cancer and dermatological diseases. *Bioorganic Medicinal Chemistry*. **14**:4323–4340.

**Nowickyj, S. M., Chithalen, J.V. and Cameron, D. (2008).** Locust retinoid X receptors: 9-Cis-retinoic acid in embryos from a primitive insect. *Proceeding National Academy of Science, U S A*. **105**:9540–9545.

**Ocaya, P., Gid'of A.C., Olofsson, P.S. and T'orm'ah, S. (2007).** CYP26 inhibitor R115866 increases retinoid signaling in intimal smooth muscle cells. *Arteriosclerosis Thrombosis and Vascular Biology* **27**:1542–1548.

**Osanai Makoto and Martin Petkovich (2005).** Expression of the Retinoic Acid-Metabolizing Enzyme CYP26A1 Limits Programmed Cell Death. *Molecular Pharmacology* **67**:1808–1817.

**Osanai, M., Sawada, N. and Lee, G. H. (2010).** Oncogenic and cell survival properties of the retinoic acid metabolizing enzyme, CYP26A1. *Oncogene* **29**:1135–1144.

**Ozpolat, B., Mehta, K., Tari, A. M. and Lopez-Berestein, G. (2002).** All-trans-retinoic acid-induced expression and regulation of retinoic acid 4-hydroxylase (CYP26) in human promyelocytic leukemia. *American Journal of Hematology* **70**:39–47.

**Ozpolat, B., Mehta, K. and Lopez-Berestein, G. (2005).** Regulation of a highly specific retinoic acid-4-hydroxylase (CYP26A1) enzyme and all-trans-retinoic acid metabolism in human intestinal, liver, endothelial, and acute promyelocytic leukemia cells. *Leukemia Lymphoma* **46**:1497–1506.

**Patrizia, D. M., Elisa M., Samantha, M., Alessandro, R., Andrea, S., Catherine, O., Maria, G., Armando, C., and Valeria, C. (2006).** Mutational screening of the *CYP26A1* gene in patients with Caudal Regression Syndrome. *Birth Defects Research (Part A)* **76**: 86–95.

**Pavez Lori`e, E., Chamcheu, J. C., Vahlquist, A. and T'orm, H. (2009).** Both all-trans retinoic acid and cytochrome P450 (CYP26) inhibitors affect the expression of vitamin A metabolizing enzymes and retinoid biomarkers in organotypic epidermis. *Archives of Dermatology Research* **301**:475–485.

**Pavez Lori`e, E., Cools, M., Borgers, M., Wouters, L. and Shroot, B. (2009).** Topical treatment with CYP26 inhibitor talarozole (R115866) dose dependently alters the expression of retinoid-regulated genes in normal human epidermis. *British Journal of Dermatology*. **160**:26–36.

**Pearce, H. L., Blanchard, K.L. and Slapak, C.A. (2008).** Failure modes in anticancer drug discovery and development. *Cancer Drug Design and Discovery. Elsevier Inc.* 424-434.

**Petkovich, M., Brand, N.J. and Krust, A. (1987).** A human retinoic acid receptor which belongs to the family of nuclear receptors. *Nature* **330**:444–50.

**Pijnappel, W. W., Hendriks, H. F., Folkers, G. E., van den Brink, C. E. and Dekker, E. J. (1993).** The retinoid ligand 4-oxo-retinoic acid is a highly active modulator of positional specification. *Nature* **366**:340–344.

**Popa, C., Dicker, A. J., Dahler, A. L. and Saunders, N. A. (1999).** Cytochrome P450, CYP26A1, is expressed at low levels in human epidermal keratinocytes and is not retinoic acid-inducible. *British Journal of Dermatology*. **141**:460–468.

**Ray, W.J., Bain, G., Yao, M. and Gottlieb, D.I. (1997).** CYP26, a novel mammalian cytochrome P450, is induced by retinoic acid and defines a new family. *Journal of Biology Chemistry*. **272**:18702–18708.

**Ren, J. H., Xiong, X. Q., Sha, Y., Yan, M.C., Lin, B., Wang, J., Jing, Y. K., Zhao, D.M. and Cheng, M. S. (2008).** Structure prediction and R115866 binding study of human CYP26A1: homology modelling, fold recognition, molecular docking and MD simulations. *Molecular Simulation* **34**:337–346.

**Rat, E., Ingrid, B.L., Delphine, A., Jean-Marc, L.G., Marie, T., Christelle, C., Nicolas, J., Isabelle, van S., Michel, L., Antonio, R., Jean-Louis, G. and Franck, B. (2006).** Evidence for a Functional Genetic Polymorphism of the Human Retinoic Acid–Metabolizing Enzyme CYP26A1, an Enzyme that may be involved in Spina Bifida. *Wiley-Liss, Inc* **76**:491–498.

**Ritchie, D. W. and Kemp, G. J. L. (2000).** Protein Docking Using Spherical Polar Fourier Correlation. *PROTEINS: Structure, Function, Genetics* **39**: 178-194.

**Roberts, A. B. and DeLuca, H. F. (1967).** Pathways of retinol and retinoic acid metabolism in the rat. *Biochemistry Journal*. **02**:600–605.

**Ross, A. C., Cifelli, C. J., Zolfaghari, R. and Li N, Q. (2011).** Multiple cytochrome P-450 genes are concomitantly regulated by vitamin A under steady-state conditions and by retinoic acid during hepatic first-pass metabolism. *Physiology Genomics* **43**:57–67.

**Ross, A.C. (1993).** Cellular metabolism and activation of retinoids: roles of cellular retinoid-binding proteins. *FASEB Journal* **7**:317–327.

**Sali, A. and Blundell, T.L. (1993).** Comparative protein modelling by satisfaction of spatial restraints. *Journal of Molecular Biology* **234**: 779-815.

**Samokyszyn, V. M., Gall, W. E. and Zawada, G. (2000).** 4-hydroxyretinoic acid, a novel substrate for human liver microsomal UDP-glucuronosyltransferase(s) and recombinant UGT2B7. *Journal of Biology Chemistry*. **275**:6908–6914.

**Schug, T.T., Berry, D.C., Shaw, N.S., Travis, S.N. and Noy. N. (2007).** Opposing effects of retinoic acid on cell growth result from alternate activation of two different nuclear receptors. *Cell*. **129**:723–733.

**Shelton, D.N., Sandoval, I.T., Eisinger, A., Chidester, S., Ratnayake, A., et al.(2006).** Up-regulation of CYP26A1 in adenomatous polyposis coli-deficient vertebrates via a WNT-dependent mechanism:implications for intestinal cell differentiation and colon tumor development. *Cancer Research*. **66**:7571–7577.

**Sippl, M. J. (1993).** Recognition of errors in three dimensional structures of proteins. *Proteins* **17**:355–362.

**Sonneveld, E., van den Brink, C.E., Tertoolen, L.G, van der Burg, B. and van der Saag, P.T. (1999).** Retinoic acid hydroxylase (CYP26) is a key enzyme in neuronal differentiation of embryonal carcinoma cells. *Developmental Biology*. **213**:390-404.

**Soprano, D.R. and Soprano, K.J.(1995).** Retinoids as teratogens. *Annual Review of Nature*. **15**:111–132.

**Stephensen, C. B. (2001).** Vitamin A, infection, and immune function. *Annual Review. Nutrition*. **21**:167–192.

**Stoppie, P., Borgers, M., Borghgraef, P., Dillen, L. and Goossens, J. (2000).** R115866 inhibits all-trans-retinoic acid metabolism and exerts retinoidal effects in rodents. *Journal of Pharmacology Experimental Theory*. **293**:304–312.

**Swindell, E. C., Thaller, C., Sockanathan, S., Petkovich, M., Jessell, T. M. and Eichele, G. (1999).** Complementary domains of retinoic acid production and degradation in the early chick embryo. *Developmental Biology*. **216**:282–296.

**Taimi, M., Helvig, C., Wisniewski, J., Ramshaw, H. and White, J. (2004).** A novel human cytochrome P450, CYP26C1, involved in metabolism of 9-cis and all-trans isomers of retinoic acid. *Journal of Biological Chemistry*. **279**:77–85.

**Tang, G.W. and Russell, R.M. (1990).** 13-cis-retinoic acid is an endogenous compound in human serum. *Journal of Lipid Research* **31**:175–182.

**Tang, X. H. and Gudas, L. J. (2011).** Retinoids, retinoic acid receptors and cancer. *Annual Reviews. Pathology*. **6**:345–364.

**Tahayato, A., Dolle, P. and Petkovich, M. (2003).** Cyp26C1 encodes a novel retinoic acid-metabolizing enzyme expressed in the hindbrain, inner ear, first brachial arch and tooth buds during murine development. *Gene Expression Patterns* **3**:449–454.

**Tay, S., Dickmann, L., Dixit, V. and Isoherranen, N. (2010).** A comparison of the roles of peroxisome proliferator activated receptor and retinoic acid receptor on CYP26 regulation. *Molecular Pharmacology*. **77**:218–227.

**Thatcher, J. E., Zelter, A. and Isoherranen, N. (2010).** The relative importance of CYP26A1 in hepatic clearance of all-trans retinoic acid. *Biochemical Pharmacology*. **80**:903–912.

**Tzimas, G. and Nau, H. (2001).** The role of metabolism and toxicokinetics in retinoid teratogenesis. *Current Pharmaceutical Design* **7**:803–831.

**Tzimas, G., Collins, M.D., Burgin, H. et al. (1996).** Embryotoxic doses of vitamin A to rabbits result in low plasma but high embryonic concentrations of all-trans-retinoic acid: risk of vitamin A exposure in humans. *Journal of Nutrition* **126**:2159–71.

**Uehara, M., Yashiro, K. and Mamiya, S. (2007).** CYP26A1 and CYP26C1 cooperatively regulate anterior posterior patterning of the developing brain and the production of migratory cranial neural crest cells in the mouse. *Developmental Biology* **302**:399–411.

**Van der Leede, B. M., van den Brink, C. E., Pijnappel, W.W., Sonneveld, E., van der Saag, P.T. and van der Burg, B. (1997).** Auto induction of retinoic acid metabolism to polar derivatives with decreased biological activity in retinoic acid-sensitive, but not in retinoic acid-resistant human breast cancer cells. *Journal of Biological Chemistry* **272**:17921–17928.

**Van Wauwe, J. P., Coene, M. C., Goossens, J., Cools, W. and Monbaliu, J. (1990).** Effects of cytochrome P-450 inhibitors on the in vivo metabolism of all-trans-retinoic acid in rats. *Journal of Pharmacology and Experimental Therapeutics* **252**:365–369.



**Van Wauwe, J. P., Coene, M. C., Goossens, J., Van Nijen, G., Cools, W. and Lauwers, W. (1988).** Ketoconazole inhibits the in vitro and in vivo metabolism of all-trans-retinoic acid). *Journal of Pharmacology and Experimental Therapeutics* **245**:718–722.

**Van Heusden, J., Van Ginckel, R., Bruwiere, H., Moelans, P. and Janssen, B. (2002).** Inhibition of all-TRANS retinoic acid metabolism by R116010 induces antitumour activity. *British Journal of Cancer* **86**:605–611.

**Vincent C. O. N., Lalji, G., Purushottamachar, P., Chopra, P., Vasaitis, T.S., Khandelwal, A., Mehta, J., Huynh, C., Belosay, A. and Patel, J. ( 2006).** Retinoic acid metabolism blocking agents (RAMBAs) for treatment of cancer and dermatological diseases. *Bioorganic & Medicinal Chemistry* **14** : 4323–4340.

**Vriend, G. (1990).** WHAT IF: a molecular modelling and drug design program. *Journal of Molecular Graph* **8**:52—56 .

**Wallace, A. C., Laskowski, R. A. and Thornton, J. M. (1995).** LIGPLOT: a program to generate schematic diagrams of protein-ligand interactions. *Protein Engineering*. **8**:127–134.

**Wang, R., Lu, Y., and Wang, S. (2003).** Comparative evaluation of 11 scoring functions for molecular docking. *Journal of Medicinal Chemistry* **46**: 2287–2303.

**Wang, Y., Zolfaghari, R. and Ross, A. C. (2002).** Cloning of rat cytochrome P450RAI (CYP26) cDNA and regulation of its gene expression by all-trans-retinoic acid *in vivo*. *Archeives of Biochemistry and Biophysics*. **401**:235–243.

**White, J. A., Boffa, M. B. and Jones, B. (1994).** A zebrafish retinoic acid receptor expressed in the regenerating caudal fin. *Development* **120**:1861–1872.

**White, J. A., Beckett-Jones, B., Guo, Y. D., Dilworth, F. J. and Bonasoro, J. (1997).** cDNA cloning of human retinoic acid-metabolizing enzyme (hP450RAI) identifies a novel family of cytochromes P450. *Journal of Biological Chemistry*. **272**:18538–18541.

**White, J. A., Guo, Y. D., Baetz, K., Beckett-Jones, B. and Bonasoro, J. (1996).** Identification of the retinoic acid-inducible all-trans-retinoic acid 4-hydroxylase. *Journal of Biological Chemistry*. **271**:29922–29927.

**White, J. A., Ramshaw, H., Taimi, M., Stangle, W. and Zhang, A. (2000).** Identification of the human cytochrome P450, P450RAI-2, which is predominantly expressed in the adult cerebellum and is responsible for all-trans-retinoic acid metabolism. *Proceeding National Academy of Science. USA* **97**:6403–6408.

**World Health Organization. (2009).** Macronutrient deficiencies: vitamin A deficiency. Available from: <http://www.who.int/nutrition/topics/vad/en/index.html>.

**Wu, L. and Ross, A. C. (2010).** Acidic retinoids synergize with vitamin A to enhance retinol uptake and STRA6, LRAT, and CYP26B1 expression in neonatal lung. *Journal of Lipid Research* **51**:378–387.

**Xi, J. and Yang, Z. (2008).** Expression of RALDHs (ALDH1As) and CYP26s in human tissues and during the neural differentiation of P19 embryonal carcinoma stem cell. *Gene Expression Patterns* **8**:438–442.

**Yadav, Madhu. and Singh, Gurmit. (2013).** Comparative Protein Structure Modeling of CytochromeP450 26A1(Cyp26A1) And Prediction of Ligand Binding Sites. *BioIT Journal of Advanced Bioinformatics Applications and Research* **4**: 30-42.

**Yadav, Madhu. and Singh, Gurmit. (2013).** BIOINFORMATICS : A New Era of Drug Design. *Indian Journal of Bioinformatics and Biotechnology* **2**:48-53.

**Yadav, Madhu. and Singh, Gurmit. (2013).** Virtual Screening of Ligand molecules for target protein CYP26A1 by using AutoDock-Vina. *International Journal of Innovative Research in Science, Engineering and Technology*. **2**:4915-4920.

**Yadav, Madhu. and Singh, Gurmit. (2013).** Docking Studies on Inhibitors for carcinogenic retinoic acid metabolizing enzyme CYP26A1. *International Journal of Advanced Research*. **1**: 117-120.

**Yamamoto, Y., Zolfaghari, R. and Ross, A. C. (2000).** Regulation of CYP26 (cytochrome P450RAI) mRNA expression and retinoic acid metabolism by retinoids and dietary vitamin A in liver of mice and rats. *FASEB Journal*. **14**:2119–2127.

**Yashiro, K., Zhao, X. and Uehara, M. (2004).** Regulation of retinoic acid distribution is required for proximodistal patterning and outgrowth of the developing mouse limb. *Developmental Cell* **6**:411–422.

**Zhang, Y., Zolfaghari, R. and Ross, A. C. (2010).** Multiple retinoic acid response elements cooperate to enhance the inducibility of CYP26A1 gene expression in liver. *Gene* **464**:32–43.

**Zhao, Q., Dobbs-McAuliffe, B. and Linney, E. (2005).** Expression of *cyp26b1* during zebrafish early development. *Gene Expression Patterns* **5**:363–369.

**Zolfaghari, R., Cifelli, C. J., Lieu, S. O., Chen, Q., Li, N. Q. and Ross, A. C. (2007).** Lipopolysaccharide opposes the induction of CYP26A1 and CYP26B1 gene expression by retinoic acid in the rat liver in vivo. *American Journal of Physiology of Gastrointestinal Liver Physiology*. **292**:1029–1036.