

LIST OF FIGURES

Figure No.		Page No.
1.1	Schematic representation of transporter (uniporter/antiporter) proteins involve in cellular Na ⁺ homeostasis and their sub-cellular localization	13
1.2	Schematic representation of salinity and drought stress signalling pathways	23
2.1	Bacterial cloning vector pGEM-T Easy	41
2.2	Schematic map of binary vector pCAMBIA1301	42
2.3	Schematic representation of T-DNA region of <i>pCAMBIA1301-SbASR-1</i> plant transformation vector construct	42
2.4	Schematic representation of promoter isolation from genomic DNA	48
2.5	Explant preparation for regeneration and transformation	56
3.1	Graphical representation of <i>SbASR-1</i> gene genomic organization	74
3.2	Representation sequences of <i>SbASR-1</i> gene and its promoter	75
3.3	Southern hybridisation of <i>S. brachiata</i> genome to determine number of paralogues of <i>SbASR-1</i> gene	79
3.4	Amino acid sequence of <i>SbASR-1</i> protein denoted in single-letter format	80
3.5	Multiple alignment of amino acid sequences of <i>SbASR-1</i> protein with other ASR-1 proteins	81
3.6	Phylogenetic tree of <i>SbASR-1</i> prepared by Maximum Likelihood method	83
3.7	Folding prediction of <i>SbASR-1</i> protein using FoldIndex program	84
3.8	Regeneration of groundnut using full and vertical halves of de-embryonated cotyledon explants via direct organogenesis	86-87
3.9	Determination of hygromycin lethal dose (LD ₅₀)	87-88
3.10	Screening of transformation efficiency using two different protocols	89
3.11	Optimization factors affecting transformation	90
3.12	Comparative analysis of GUS transient expression efficiency and subsequent regeneration efficiency on SM3 media	92
3.13	Genetic transformation, regeneration, selection, hardening and seed development of putative transgenic lines	95

3.14	Flow chart showing optimized transformation and regeneration protocol for groundnut cv. GG 20	96
3.15	PCR confirmation of putative transgenic lines	97
3.16	Southern blotting and hybridization of T0 putative transgenic lines	98
3.17	Reverse transcriptase PCR of <i>SbASR-1</i> gene in transgenic lines	99
3.18	Histochemical GUS assay: of leaves from Wt (a) and putative transgenic lines (b & c)	99
3.19	Histochemical GUS assay of Wt and transgenic lines in T1 generation	100
3.20	Morphological differences in Wt and transgenic lines at control and stress treatment conditions	101
3.21	Estimation of photosynthetic pigments	103-104
3.22	Estimation of electrolyte leakage (a), proline (b), and RWC (c) from leaves of Wt plants and transgenic lines under control and stress conditions	106
3.23	Estimation of MDA content to determine the lipid peroxidation under stress	107
3.24	Estimation of total soluble sugar (a), reducing sugar (b), and starch content in Wt plants and transgenic lines	108-109
3.25	DAB and NBT staining of Wt plants leaves and transgenic lines leaves	110
3.26	Relative fold change in transcript expression of <i>APX</i> (a), <i>CAT</i> (b) and <i>SOD</i> (c) compared to control plants	112