

LIST OF TABLES

TABLE NO.	TITLE	PAGE NO.
1.1	Factors and parameters influencing bacterial mineral oxidation and metal mobilisation	5
1.2	Differentiation tests of coliform bacilli	31
1.3	Tests to identify the species level in <i>Serratia</i>	32
3.1a	Amino acid bound to the copper atom forming the functional active site in the different blue copper proteins of eukaryotic and prokaryotic origin	87
3.1b	Number of identities for each protein from different databases	87
3.1c	Existing signatures for the different blue copper proteins	88
3.1d	Number of identities the already existing signatures brings out from the different non redundant databases	90
3.1e	Different signatures designed based on the functional homologous region of each of the blue copper protein	91
3.2a	Microbiological assays carried out for the identification of the soil isolate <i>Serratia</i> sp	93
3.2b	Important biochemical tests for identification of soil isolate <i>Serratia</i> sp	93
3.3a	Comparative analysis of prodigiosin production by <i>Serratia marsescens</i> in different media at 28 ⁰ C, 30 ⁰ C and 37 ⁰ C temperatures	100

TABLE NO.	TITLE	PAGE NO.
3.3b	Comparative analysis of extracellular protein production by <i>Serratia marsescens</i> in different media at 28 ⁰ C, 30 ⁰ C and 37 ⁰ C temperatures	100
3.4	Comparative analysis of pigment production	101
3.5a	Comparative analysis of prodigiosin production by <i>Serratia marsescens</i> in different seed media at 28 ⁰ C, 30 ⁰ C and 37 ⁰ C temperatures	103
3.5b	Comparative analysis of extracellular protein production by <i>Serratia marsescens</i> in different seed media at 28 ⁰ C, 30 ⁰ C and 37 ⁰ C temperatures	103
3.6	Comparative analysis of sesame broth, sesame oil broth, peanut seed broth, peanut oil broth, copra seed broth and coconut oil broth	105
3.7	Concentration of prodigiosin in mg/ml extracted from different pH conditions	107
3.8	Extracellular protein and pigment produced by <i>Serratia marsescens</i> in peanut medium in the presence of different metals of standard concentration of 0.2 mg/ ml.	113
3.9	Concentration of intracellular and extracellular protein expressed in mg/ml during the growth of <i>Serratia marsescens</i> in peanut medium in the presence of the four metals.	122
3.10	<i>Serratia marsescens</i> cultures which had grown in the presence of nickel, copper, cobalt and chromium and had a block in the pigment production, when reverted back to medium without metal in it had shown pigment production	122