

## TABLE OF CONTENTS

Chapter No	CONTENT	Page no
<b>1</b>	<b>INTRODUCTION</b>	<b>1</b>
1.1	Liquid Crystals	1
1.2	Classification of liquid crystals	2
1.2.1	Nematic liquid crystals	2
1.2.2	Smectic liquid crystals	4
1.2.3	Compounds exhibiting only the nematic liquid crystalline phase	6
1.2.4	Compounds exhibiting the smectic A liquid crystalline phase	7
1.3	Theories of phase transitions	8.
1.3.1	Landau theory of phase transitions	8
1.3.2	Landau- de- Gennes theory of the N-I transition	10
1.3.3	Landau-de -Gennes theory for the N - SmA transition	11
<b>2.</b>	<b>LITERATURE SURVEY AND OBJECTIVE</b>	<b>14</b>
2.1	Review of earlier experimental work	14
2.2	Review of earlier theoretical work	18
2.2.1	Considering the effect of surface orientation potential	18
2.2.2	Considering the Quenching of director fluctuations in thin cells	20
2.2.3	Combined effect	20
2.3	Motivation and Objective of the present work	23
2.4	Importance of the study	24

<b>3.</b>	<b>EXTENSION OF M-S THEORY</b>	<b>25</b>
3.1	Review of Maier-Saupe theory for the N-I transition	25
3.2	Extension of M-S Theory for nematic liquid crystals in thin cells.	27
3.2.1	Theoretical Model	27
3.2.2	Expression for Free energy and order parameter	28
3.2.3	Method of Calculation	30
3.2.4	Results and Discussion.	31
3.2.5	Conclusions:	36
<b>4.</b>	<b>A MOLECULAR THEORY FOR PHASE DIAGRAMS INVOLVING SMECTIC AND NEMATIC PHASES</b>	<b>37</b>
4.1	Introduction	37
4.2	The Sm A-N Transition	38
4.3	Theory of the Smectic A –Nematic transition	38
4.4	Review of the McMillan theory of SmA liquid crystals	39
4.5	Extension of Mc-Millan Theory for thin cell.	41
4.5.1	Theoretical Model	42
4.5.2	Expression for Free energy and order parameter	43
4.5.3	Method of Calculation	45
4.5.4	Results and Discussion.	46
4.5.5	Conclusions:	54
<b>5.</b>	<b>A MOLECULAR THEORY OF THE NEMATIC TO NEMATIC PHASE TRANSITION IN STRONGLY POLAR COMPOUNDS.</b>	<b>55</b>
5.1	Introduction	55

5.2	Model for molecular pairs with $\parallel$ dipole moments at low temperatures	58
5.3	Polar liquid crystals in thin cells.	62
5.4	Extension of the polar model to liquid crystals in thin cells.	63
5.4.1	Assumptions	63
5.4.2	Free energy and order parameters	65
5.4.3	Calculations	67
5.4.4	Results and Discussion.	68
5.4.5	Conclusions	72
<b>6.</b>	<b>REFERENCES</b>	

## LIST OF TABLES

TABLE	PAGE
	NO
TABLE 2.1 Comparison of $\Delta n$ in thin and thick cells	17
Table 2.2 Comparison of calculated values of the order parameters in $1\mu\text{m}$ using different approximations and the bulk values at a few temperatures.	22
Table 3.1: Comparison of calculated values of order parameters (S) in cells of thicknesses $1\mu\text{m}$ and $10\mu\text{m}$ for first potential (exponential decay)	34
Table 3.2: Comparison of calculated values of order parameters (S) in cells of thicknesses $1\mu\text{m}$ and $10\mu\text{m}$ for the second potential (inverse square type)	34
Table 3.3: Comparison of $dS\%$ in first and second potential	35
Table 4.1: Comparison of calculated values of average order parameters (S) in cells of thicknesses $1\mu\text{m}$ and $10\mu\text{m}$ with nematic phase only	

Table 4.2: Comparison of calculated values of average order parameters ( $S$ ) in cells of thicknesses $1\mu\text{m}$ and $10\mu\text{m}$ with smectic near the surface.	52
Table 4. 3: Comparison of $dS$ % in first and second cases	53
Table 5.1: Comparison of calculated values of order parameters ( $S$ ) in cells of thicknesses $1\mu\text{m}$ and $10\mu\text{m}$ .	71

## LIST OF FIGURES

FIGURE	PAGE NO
Fig. 1.1. Schematic representation of the ordering of the rod-like molecules in the isotropic and the nematic phases. The director $\hat{n}$ and the choice of axes in the laboratory frame of reference are also shown.	2
Fig. 1.2. Schematic representation of ordering of rod-like molecules in the smectic A and the smectic C phases, showing the nematic director $\hat{n}$ , the average spacing $d$ between the layers, the wave vector $\vec{q}$ representing the density wave along the layer normal. $\vec{q}$ and $\hat{n}$ are parallel in the smectic A liquid crystals whereas in the smectic C liquid crystals, $\hat{n}$ is tilted with respect to $\vec{q}$ by an angle $\omega$ .	5
Fig. 2.1. Transmitted intensity profile as a function of temperature for cells of thickness (a) $3\mu\text{m}$ and (b) $1.9\mu\text{m}$ respectively.	15
Fig.2.2. $\Delta n$ with temperature for the compound CP7B o: cell of thickness $\sim 1.5\mu\text{m}$ and +: cell of thickness $\sim 25\mu\text{m}$ .	15

Fig.2.3(a). The variations of birefringence ( $\Delta n \propto S$ ) of CN as a function of temperature are shown for cells of 3 different thickness: 1.5 $\mu\text{m}$ (open circles) 2.2 $\mu\text{m}$ (open triangles) and 14 $\mu\text{m}$ (open squares).	
Fig.2.3(b). Variations of $\Delta n$ as functions of temperature of the mixture of CN and BAN in three different cell thicknesses, namely 1.4 $\mu\text{m}$ (open circles).	17
Fig.2.4. Cross section of a sample cell. Hatched region is SiO evaporated film: $n_o$ , $n$ and $\bar{n}$ are the refractive indices of the cell plates, bulk of LC, and LC near the surface, respectively	18
Fig 2.5. Temperature dependence of order parameters in the bulk and at the surface	18
Fig. 2.6. Nematic order near a free surface of the bulk isotropic phase	19
Fig. 3.1. Variation of order parameter ( $S_z$ ) with distance from surface ( $z$ ) for a cell of thickness 1 $\mu\text{m}$ , upto middle of cell, with $r_0 = 0.01 \mu\text{m}$ , at $T = T_{NI} - 1^0$ and $T_{NI} - 10^0$	32
Fig. 3.2. Variation of order parameter ( $S_z$ ) with distance from surface ( $z$ ) for a cell of thickness 10 $\mu\text{m}$ , upto middle of cell, with $r_0 = 0.01 \mu\text{m}$ , at $T = T_{NI} - 1^0$ and $T_{NI} - 10^0$	33
Fig. 4.1 Phase diagram for theoretical model parameter $\alpha$ ; Inset: typical phase diagram for homologues series of compounds showing transition temperatures versus length of the alkyl end-chains	41
Fig. 4.2. Calculated phase diagram as a function of $T_r$ and $\alpha$ without considering the surface potential. Arrow indicates $\alpha = 0.95$ which is chosen for the rest of the calculations.	47
Fig. 4.3. The calculated variation of order parameter ( $S_z$ ) with	

distance from surface ( $z$ ) for a cell of thickness  $1\mu\text{m}$ , with  $r_0 = 0.01\mu\text{m}$ , at  $T = T_{\text{NI}} - 1^\circ$ , with  $\alpha = 0.95$ . The arrow indicates  $z/r_0 = 4$  at which smectic becomes nematic. The calculated variation of  $S_Z$  with  $z$  considering only the nematic phase ( chapter- 3 ) is also shown for comparison

48

Fig.4.4. Calculated distance-temperature phase diagram showing smectic (Sm), nematic (N), para nematic(Np) and isotropic phases. The Np phase becomes isotropic for large values of  $z/r_0$  (not shown)

49

Fig.4.5. Calculated variation of nematic and smectic order parameters as a function of  $z/r_0$  showing the presence of smectic phase upto  $z/r_0 = 5$ , for  $\alpha = 0.95$  and  $T = T_{\text{NI}} - 10^\circ$  corresponding to the phase diagram in figure 4.4

49

Fig.4.6. Variation of  $S_Z$  with  $z/r_0$  at different  $T_r$  corresponding to nematic to paranematic transitions shown in the phase diagram in figure 4.4. Solid lines from right to left correspond to  $T_r = 1.03$  to  $1.08$ . The dashed line shows decrease in  $S_Z$  as  $T_r$  is increased

50

Fig. 5.1. (a) Variation  $C_p$  for a sample with  $X = 53.7\%$ , where  $X$  is the mole percent concentration of  $\text{DB}_{10}\text{ONO}_2$  in  $\text{DB}_8\text{ONO}_2$ . The strong peak in the middle corresponds to the  $N_d-N_1$  transition and the small peak marked by the right arrow corresponds to  $\text{SmA}_d-N_d$  transition and the left arrow to  $\text{SmA}_1-N_1$  transition. (b) Detail of the  $T-X$  phase diagram for the same mixture for  $X$  between  $51\%$  and  $55\%$ . Filled circles, open circles and open squares represent calorimetric, X-ray scattering and optical measurements respectively. The dashed curves indicate  $2^{\text{nd}}$  order phase transitions and the solid curves,  $1^{\text{st}}$  order ones. TCP marks the tricritical point

57



Figure-5.2. Variation in the transmitted intensity as a function of temperature for the compound CP7B for cell thickness of  $1.9 \mu\text{m}$ . ( Note the jump around  $33^\circ \text{C}$ ). 57

Figure-5.3. General phase diagrams predicted by the dislocation loop melting theory of Prost and Toner.  $A_1$  and  $A_2$  are general variables which can be mapped to pressure and temperature or pressure and concentration etc 58

Fig. 5.4. Schematic diagram showing (a) the antiparallel configuration of two molecules favoured at intermediate molecular separations and (b) the parallel configuration favoured at relatively low values of intermolecular separation. The arrow with a solid line represents the permanent dipole moment and the one with a dotted line, the induced one. (For the sake of clarity, the relative separation in (a) is exaggerated) 59

Fig. 5.5. The pairing energy difference  $\Delta E = E_A - E_P$  (in  $10^{-19}\text{J}$ ) between the antiparallel and parallel configurations plotted as a function of the intermolecular separation  $r$  (in  $\text{\AA}$ ) 61

Fig. 5.6. Calculated variation of  $S_{av}$  and  $X_P$  as a function of distance from the cell wall ( $z/r_0$ ). In the legend, '1' corresponds to  $T = T_{NI} - 0.2^\circ$  and '2' to  $T = T_{NI} - 10^\circ$  69

Figure-5.7. Calculated variation of weighted order parameters  $X_A S_A$  and  $X_P S_P$  and the average order parameter  $S_{av}$  as a function of distance from the cell wall ( $z/r_0$ ) 69