

## LIST OF FIGURES

Figure No.	TITLE	Page No.
1	Possible mechanism involved in the genesis of PDN	39
2	Possible involvement of peroxynitrite (ONOO-) in PDN	86
3	Standard curve for tissue TBARS level using Malondialdehyde	100
4	Standard curve for reduced glutathione using Glutathione (GSH)	100
5	Standard curve for tissue nitrite/nitrate level using Sodium nitrite	101
6	Standard curve for H <sub>2</sub> O <sub>2</sub> using H <sub>2</sub> O <sub>2</sub>	101
7	Standard curve for tissue catalase level using catalase	102
8	Effect of STZ-induced hyperglycemia on pain sensitivity to thermal test (Hot-Plate –Test) in mice	116
9	Effect of STZ-induced hyperglycemia on pain sensitivity to thermal test (radiant heat) (Planter-test) in mice	116
10	Effect of STZ-induced hyperglycemia on pain sensitivity to thermal test (radiant heat-Tail-Flick Test) in Mice	117
11	Effect of STZ-induced hyperglycemia on pain sensitivity to mechanical pressure-test in mice	117
12	Effect of STZ-induced hyperglycemia on pain sensitivity to cold-test (water) (allodynia) in Mice	118
13	Effect of STZ-induced hyperglycemia on pain sensitivity to Cold (Acetone) -test in mice	118
14	Effect of STZ-induced hyperglycemia on Carrageenan induced pain sensitivity in mice	119
15	Effect of STZ-induced hyperglycemia on formalin induced flinching behaviors in mice	119
16	Effect of STZ-induced hyperglycemia on sensitivity to thermal (Tail-Flick Test) stimuli on antinociceptive effect of opioids	120
17	Effect of STZ-induced hyperglycemia on the sensitivity to antinociceptive effect of morphine in thermal test.	120
18	Effect of STZ-induced experimental diabetes on the sensitivity to antinociceptive effect of morphine in pressure	121

	test	
<b>19</b>	Effect of STZ-induced hyperglycemia on the sensitivity to antinociceptive effect of DAMGO in thermal test	121
<b>20</b>	Effect of STZ-induced Experimental Diabetes On The Sensitivity to Antinociceptive Effect of DAMGO In Pressure Test	122
<b>21</b>	Effect of STZ-induced experimental diabetes on oxycodone and DPDPE induced % MPE in Thermal Tes	122
<b>22</b>	Effect of SHS-treatment of non diabetic mice on antinociceptive effect of morphine and DAMGO	123
<b>23</b>	Effect of STZ-induced hyperglycemia and Splenectomy and SHS on antinociceptive effect of Morphine	123
<b>24</b>	Effect of STZ-induced hyperglycemia, splenectomy and SHS on antinociceptive effect of DAMGO in mice	124
<b>25</b>	Effect of STZ-induced hyperglycemia and SHS on antinociceptive effect of DPDPE and Oxycodone in mice	124
<b>26</b>	Effect of STZ-induced hyperglycemia on the sensitivity to antinociceptive effect of WIN 55,212-2 in mice	125
<b>27</b>	Effect of STZ-induced hyperglycemia on the Sensitivity to mechanical stimuli on antinociceptive effect of WIN 55,212-2 in mice	125
<b>28</b>	Effect of STZ-induced hyperglycemia and pharmacological intervention on antinociceptive effect of cannabinoids	126
<b>29</b>	Effect of SHS on antinoceptive effect of cannabis extract and Win 55, 212-2	126
<b>30</b>	Effect of STZ-induced hyperglycemia and pharmacological intervention on the sensitivity to antinociceptive effect of NSAIDs	127
<b>31</b>	Effect of STZ-induced hyperglycemia and pharmacological interventions on antinociceptive effect of neurosteroids	127
<b>32</b>	Effect of testosterone, progesterone, DHEA and allopregnlone on carrageenan induced inflammatory hyperalgesia	128
<b>33</b>	Effect of Opioid antagonist on STZ-induced Antinociceptive Tolerance of Morphine and DAMGO In Mice	128
<b>34</b>	Effect of opioid antagonist on STZ-induced antinociceptive tolerance to WIN 55,212 in mice	129
<b>35</b>	Effect of low dose of rimonabant on STZ-induced antinociceptive tolerance to WIN 55,212-2 in mice	129
<b>36</b>	Effect of low dose of cannabinoid- antagonist rimonabant (RIM) on STZ-induced morphine and DAMGO analgesic	130

	tolerance in mice	
<b>37</b>	Effect of Cyclosporine-A, Thalidomide, Pentoxyphylline and Minocycline on the Development of thermal hyperalgesia (tail-flick test) in STZ-treated Mice	130
<b>38</b>	Effect of Cyclosporine-A, Thalidomide, Pentoxyphylline and Minocycline on the Development of mechanical allodynia in STZ- treated Mice	131
<b>39</b>	Effect of Cyclosporine-A, Thalidomide, Minocycline and Pentoxyphylline on Established Hyperalgesia in Diabetic Mice	131
<b>40</b>	Effect of Cyclosporine-A, Thalidomide, Minocycline and Pentoxyphylline on Established Allodynia in Diabetic Mice	132
<b>41</b>	Effect of Thalidomide and Pentoxyphyllne on STZ-induced in Decreased Antinociceptive effect of Morphine and DAMGO	132
<b>42</b>	Effect of Thalidomide and Pentoxyphylline on STZ-induced- Decreased Antinociceptive effect of Cannabis extract and Win 55,212-2	133
<b>43</b>	Effect of Thalidomide and Pentoxyphyllne on STZ-induced Decreased in Antinociceptive effect of NSAIDs	133
<b>44</b>	Effect of Thalidomide and Pentoxyphyllne on STZ-induced Decreased in Antinociceptive effect of Neurosteroids	134
<b>45</b>	Effect of Cyclosporine-A and Minocycline on STZ-induced Decrease in Antinociceptive effect of Morphine and DAMGO	134
<b>46</b>	Effect of Cyclosporine-A and Minocycline on STZ-induced Decreased in Antinociceptive effect of Cannabinoids	135
<b>47</b>	Effect of Cyclosporine –A and Minocycline on STZ-induced Decreased in Antinociceptive effect of NSAIDs	135
<b>48</b>	Effect of Cyclosporine-A and Minocycline on STZ-induced Decreased in Antinociceptive effect of Neurosteroids	136
<b>49</b>	Effect of Dextromethorphan and Ketamine on STZ-induced decrease in % MPE of Morphine and DAMGO	136
<b>50</b>	Effect of Dextromethorphan and Ketamine on STZ-induced decrease in % MPE of Cannabis extract and WIN 55,212-2	137
<b>51 (a)</b>	Effect of Dextromethorphan on STZ-induced decrease in % MPE of NSAIDs	137
<b>51 (b)</b>	Effect of Ketamine on STZ-induced decrease in % MPE of NSAIDs	138
<b>52</b>	Effect of Dextromethorphan and Ketamine on STZ-induced decrease in % MPE of Neurosteroids	138
<b>53</b>	Effect of chelerythrine on Hyperglycemia- induced decrease in % MPE of opioids	139
<b>54</b>	Effect of chelerythrine on Hyperglycemia- induced	139

	decrease in % MPE of cannabinoids	
<b>55</b>	Effect of L-NAME and Aminoguanidine, an iNOS inhibitor, on the Development of STZ-induced thermal hyperalgesia	140
<b>56</b>	Effect of Aminoguanidine and L-NAME on the Development of STZ-induced mechanical allodynia	140
<b>57</b>	Effect of Aminoguanidine and L-NAME on Established (STZ-induced) Thermal Hyperalgesia	141
<b>58</b>	Effect of Aminoguanidine and L-NAME on Established (STZ-induced) Mechanical allodynia	141
<b>59</b>	Effect of Aminoguanidine and L-NAME on Hyperglycemia- induced reduced sensitivity of Opioids (% MPE)	142
<b>60</b>	Effect of Aminoguanidine and L-NAME on Hyperglycemia- induced decrease in % MPE of Cannabinoids	142
<b>61</b>	Effect of Aminoguanidine and L-NAME on Hyperglycemia- induced decrease in % MPE of NSAIDs	143
<b>62</b>	Effect of Aminoguanidine and L-NAME on Hyperglycemia- induced decrease in % MPE of neuroseroids	143