

LIST OF FIGURES

Figure No.	Title
1.1	Sampling Areas
1.2	Isolates obtained from different mangrove sediments samples
1.3	Primary screening for selected mangrove actinomycetes
1.4	Antimicrobial activity of different concentration of supernatant using secondary screening methods (+17 actinomycetes)
1.5	Antimicrobial activity of different concentration of supernatant using secondary screening methods (+18 actinomycetes)
1.6	Antimicrobial activity of different concentration of supernatant using secondary screening methods (+50 actinomycetes)
1.7	Antimicrobial activity of different concentration of supernatant using secondary screening methods (+100 actinomycetes)
1.8	Antimicrobial activity of different concentration of supernatant using secondary screening methods (+112 actinomycetes)
1.9	Antimicrobial activity of different concentration of supernatant using secondary screening methods (+118 actinomycetes)
1.10	16s rRNA gene, Partial sequencing of actinomycetes +18 strain
1.11	16s rRNA gene, Partial sequencing of actinomycetes +50 strain
1.12	16s rRNA gene, Partial sequencing of actinomycetes +118 strain
1.13	16s rDNA gene sequencing of actinomycetes strain +18 by Maximum likelihood tree methods
1.14	16s rDNA gene sequencing of actinomycetes strain +50 by Maximum likelihood tree methods
1.15	16s rDNA gene sequencing of actinomycetes strain +118 by Maximum likelihood tree methods
1.16	Predicted structure of 16s rDNA of actinomycetes strain +18
1.17	Predicted structure of 16s rDNA of actinomycetes strain +50
1.18	Predicted structure of 16s rDNA of actinomycetes strain +118
2.1	UV-Vis spectrum of bioactive compounds from actinomycetes +18 strain
2.2	UV-Vis spectrum of bioactive compounds from actinomycetes +50 strain
2.3	FTIR spectrum of bioactive compounds from actinomycetes +18 strain
2.4	FTIR spectrum of bioactive compounds from actinomycetes +50 strain
2.5	NMR spectrum of bioactive compounds from actinomycetes +18 strain
2.6	NMR spectrum of bioactive compounds from actinomycetes +50 strain
	UV- Visible spectra of actinomycete supernatant (+18) filtrate as a function of time at different incubations with silver nitrate (1mM) after 24h, 48h and 72h. respectively. The peak 420nm corresponds to the plasmon resonance of silver nanoparticles.
3.2	UV- Visible spectra of actinomycetes supernatant (+50) filtrate as a function of time at different incubations with silver nitrate (1mM) after 24h, 48h and 72h. respectively. The peak 420nm corresponds to the plasmon resonance of silver nanoparticles.
3.3	UV- Visible spectra of actinomycetes supernatant (+118) filtrate as a function of time at different incubations with silver nitrate (1mM) after 24h, 48h and 72h. respectively. The peak 420nm corresponds to the plasmon resonance of silver nanoparticles.
3.4	XRD spectra of silver nanoparticles synthesized by +18 actinomycetes
3.5	XRD spectra of silver nanoparticles synthesized by +50 actinomycetes

3.6	XRD spectra of silver nanoparticles synthesized by +118 actinomycetes
3.7	FTIR Spectrum of vacuum dried powder of silver nanoparticles synthesis from 1mM AgNO ₃ and actinomycetes +18 supernatant
3.8	FTIR Spectrum of vacuum dried powder of silver nanoparticles synthesis from 1mM AgNO ₃ and actinomycetes +50 supernatant
3.9	FTIR Spectrum of vacuum dried powder of silver nanoparticles synthesis from 1mM AgNO ₃ and actinomycetes +118 supernatant
4.1	Length of <i>P. monodon</i> fed with different concentrations of actinomycetes probiotics
4.2	Weight gain of <i>P. monodon</i> fed with different concentrations of actinomycetes probiotics
4.3	Survival rate of <i>P. monodon</i> fed with different concentrations of actinomycetes probiotics
4.4	Total Heterotrophic bacterial counts in experimental and control tanks
4.5	Total <i>Vibrio</i> counts in experimental and control tanks