

## List of Tables

Table No.	Title	Page No.
Table 2.1	Physical data of 3,6-dihydro-2 <i>H</i> -pyrans ( <b>3</b> ).	46
Table 2.2	IR, <sup>1</sup> H NMR and <sup>13</sup> C NMR data of 3,6-dihydro-2 <i>H</i> -pyrans ( <b>3</b> ).	48
Table 2.3	Computed HDA reactions of aromatic aldehydes with 1,3-dienes in the absence and presence of TiCl <sub>4</sub> (Scheme 2.2).	61
Table 2.4	Selected bond lengths (Å) and their Wiberg Bond Indices (in parentheses) of the transition structures in the model HDA reactions 1-19.	67
Table 2.5	Selected bond lengths (Å) and their Wiberg Bond Indices (in parentheses) of the intermediates and products in the model HDA reactions 1-19.	69
Table 2.6	NICS(0) and NICS(1) values calculated for the transition structures involved in the TiCl <sub>4</sub> catalyzed HDA reaction of aldehydes with isoprene having O/Me, 1:3 orientation along with uncatalyzed HDA reactions.	71
Table 2.7	Total energies of different species and the relative enthalpies, Δ <i>H</i> for the HDA reactions 1-19.	72
Table 2.8	Calculated ratios of the regioisomers for the HDA reactions between <b>4</b> and <b>2b</b> .	76
Table 2.9	Global properties of the dienes ( <b>2</b> ), aromatic aldehydes ( <b>4a-c</b> ) and their complexes with TiCl <sub>4</sub> ( <b>4d-f</b> ).	77
Table 2.10	Stabilization energies <i>E<sub>ij</sub></i> from second-order perturbation NBO analysis of the transition structures and intermediate of the HDA reaction 10 at the DFT (B3LYP/6-31G(d,p)) level.	79
Table 3.1	Physical data of 3,6-dihydro-4,5-dimethyl-2 <i>H</i> -pyrans ( <b>4</b> ).	88
Table 3.2	IR, <sup>1</sup> H NMR and <sup>13</sup> C NMR data of 3,6-dihydro-4,5-dimethyl-2 <i>H</i> -pyrans ( <b>4</b> ).	89
Table 3.3	HPLC data of 3,6-dihydro-4,5-dimethyl-2 <i>H</i> -pyrans ( <b>4</b> ) using chiral OJ-H column.	96
Table 3.4	Comparison of the enthalpies of the DA reactions of aromatic aldehydes with DMB in the absence/presence of TiCl <sub>4</sub> as catalyst at different levels (Scheme (3.2)).	101
Table 3.5	Selected bond lengths (Å) and their Wiberg Bond Indices (in parentheses) of the reactants ( <b>5</b> , <b>2</b> ), transition structures ( <b>TS1-TS4</b> ) and cycloadducts ( <b>6a-6d</b> ) for the model reactions 1-4.	107

---

Table 3.6	Selected bond angles and dihedral angles (in degrees) of the transition structures ( <b>TS1-TS4</b> ), and the corresponding cycloadducts ( <b>6a-6d</b> ) for the model reactions 1-4.	108
Table 3.7	Total energies and relative enthalpies ( $\Delta H$ ) of the reactants, cycloadducts and transition structures in the gas phase and in toluene for the model reactions 1-4.	109
Table 3.8	Standard state entropies, $S^\circ$ , entropy change, $\Delta S$ , relative enthalpies, $\Delta H$ and relative Gibbs free energies $\Delta G$	111
Table 3.9	Ratio of the stereoisomers in the gas and toluene phases for the asymmetric HDA reaction in the <i>endo</i> manner.	112
Table 4.1	Investigated HDA reactions of benzaldehyde with 1,3-diene (Scheme 4.1).	117
Table 4.2	Selected bond lengths ( $\text{\AA}$ ) and their WBI of the transition structures for the model HDA reactions 1-12.	125
Table 4.3	Selected bond lengths ( $\text{\AA}$ ) and their WBI of the products for the model HDA reactions 1-12.	126
Table 4.4	NICS(0) and NICS(1) values calculated for the transition structures with <i>endo</i> approach.	128
Table 4.5	Total energies of different species and relative enthalpies for the HDA reactions 1-12.	128
Table 4.6	Calculated ratios of the regioisomers for the HDA reactions between <b>1</b> and <b>2b</b> .	131
Table 4.7	FMO's ( $E_{\text{HOMO}}$ and $E_{\text{LUMO}}$ ) energies, electronic chemical potential $\mu$ , chemical hardness $\eta$ , global electrophilicity $\omega$ and global nucleophilicity $N$ of the Lewis acid complexes of benzaldehyde ( <b>1a-d</b> ) and dienes ( <b>2</b> ).	132
Table 4.8.	Physical data of 3,6-dihydro-4,5-dimethyl-2 <i>H</i> -pyrans ( <b>5</b> )	134
Table 4.9.	IR, $^1\text{H}$ NMR and $^{13}\text{C}$ NMR data of 3,6-dihydro-4,5-dimethyl-2 <i>H</i> -pyrans ( <b>5</b> )	135

---

## List of Figures

Figure No.	Title	Page No.
Figure 1.1	DA reaction of conjugated dienes with dienophiles.	1
Figure 1.2	NED and IED HDA reactions involving carbonyl functionality.	2
Figure 1.3	FMO presentation showing the effect of Lewis acids on carbonyl groups.	3
Figure 1.4	Possible pathways for the HDA reaction of carbonyl compound.	3
Figure 1.5	Concerted and stepwise cycloaddition mechanisms for the HDA reaction of carbonyl compounds.	4
Figure 1.6	Activation of carbonyl group by a hydrogen bond interaction with a protic organic solvent	6
Figure 2.1	IR spectrum of 3,6-dihydro-4,5-dimethyl-2-(3'-nitrophenyl)-2 <i>H</i> -pyran ( <b>3a</b> ).	53
Figure 2.2	<sup>1</sup> H NMR spectrum of 3,6-dihydro-4,5-dimethyl-2-(3'-nitrophenyl)-2 <i>H</i> -pyran ( <b>3a</b> )	55
Figure 2.3	<sup>1</sup> H NMR spectrum of 3,6-dihydro-4-methyl-2-(3'-nitrophenyl)-2 <i>H</i> -pyran ( <b>3g</b> )	56
Figure 2.4	<sup>13</sup> C NMR spectrum of 3,6-dihydro-4,5-dimethyl-2-(3'-nitrophenyl)-2 <i>H</i> -pyran ( <b>3a</b> )	58
Figure 2.5	<sup>13</sup> C NMR spectrum of 3,6-dihydro-4-methyl-2-(3'-nitrophenyl)-2 <i>H</i> -pyran ( <b>3g</b> )	59
Figure 2.6	Optimized geometries of the stationary points in the HDA reaction of <b>4a</b> with <b>2a</b> in the <i>endo</i> and <i>exo</i> manners.	63
Figure 2.7	Optimized geometries of the stationary points in the HDA reaction of <b>4a</b> with <b>2b</b> in the <i>endo</i> manner.	64
Figure 2.8	Optimized geometries of the stationary points in the HDA reaction of <b>4d</b> with <b>2a</b> in the <i>endo</i> manner.	65
Figure 2.9	Optimized geometries of the stationary points in the HDA reaction of <b>4d</b> with <b>2b</b> in the <i>endo</i> manner.	65

Figure No.	Title	Page No.
Figure 2.10	Energy profile diagram for the reaction of benzaldehyde and DMB in the absence and presence of TiCl <sub>4</sub> .	74
Figure 2.11	LUMOs of benzaldehyde ( <b>4a</b> ) and complexes of aromatic aldehydes with TiCl <sub>4</sub> ( <b>4d-f</b> ) and the HOMO of DMB ( <b>2a</b> ), calculated at the B3LYP/6-31G(d,p) level.	80
Figure 3.1	IR spectrum of 3,6-dihydro-4,5-dimethyl-2-(2'-methoxyphenyl)-2 <i>H</i> -pyran ( <b>4a</b> )	92
Figure 3.2	<sup>1</sup> H NMR spectrum of 3,6-dihydro-4,5-dimethyl-2-(2'-methoxyphenyl)-2 <i>H</i> -pyran ( <b>4a</b> )	93
Figure 3.3	<sup>13</sup> C NMR spectrum of 3,6-dihydro-4,5-dimethyl-2-(2'-methoxyphenyl)-2 <i>H</i> -pyran ( <b>4a</b> )	95
Figure 3.4	Chiral HPLC spectrum of 3,6-dihydro-4,5-dimethyl-2-(2'-methoxyphenyl)-2 <i>H</i> -pyran ( <b>4a</b> )	97
Figure 3.5	Chiral HPLC spectrum of 3,6-dihydro-4,5-dimethyl-2-(3'-nitrophenyl)-2 <i>H</i> -pyran ( <b>4b</b> )	98
Figure 3.6	Chiral HPLC spectrum of 3,6-dihydro-4,5-dimethyl-2-(3'-trifluoromethylphenyl)-2 <i>H</i> -pyran ( <b>4c</b> )	99
Figure 3.7	Chiral HPLC spectrum of 3,6-dihydro-4,5-dimethyl-2-(4'-trifluoromethylphenyl)-2 <i>H</i> -pyran ( <b>4d</b> )	100
Figure 3.8	Optimized geometries of all the stationary points for the HDA reaction of <b>5</b> with <b>2</b> from <i>Si</i> face in the <i>endo</i> and <i>exo</i> manner.	105
Figure 3.9	Optimized geometries of all the stationary points for the HDA reaction of <b>5</b> with <b>2</b> from <i>Re</i> face in the <i>endo</i> and <i>exo</i> manner.	106
Figure 3.10	Energy profiles of the attack of DMB ( <b>2</b> ) on <i>Si</i> and <i>Re</i> faces of the >C=O functionality of benzaldehyde complex ( <b>5</b> ) calculated at the B3LYP/STO-3G level in gas phase and in toluene.	110
Figure 4.1	Optimized geometries of the stationary points on the PES of the HDA reaction of <b>1a</b> with <b>2a</b> in the <i>endo</i> manner.	119
Figure 4.2	Optimized geometries of the stationary points on the PES of the HDA reaction of <b>1b</b> with <b>2a</b> in the <i>endo</i> manner.	119
Figure 4.3	Optimized geometries of the stationary points on the PES of the HDA reaction of <b>1c</b> with <b>2a</b> in the <i>endo</i> manner.	120

Figure No.	Title	Page No.
Figure 4.4	Optimized geometries of the stationary points on the PES of the HDA reaction of <b>1d</b> with <b>2a</b> in the <i>endo</i> manner.	120
Figure 4.5	Optimized geometries of the stationary points on the PES of the HDA reaction of <b>1a</b> with <b>2b</b> in the <i>endo</i> manner.	121
Figure 4.6	Optimized geometries of the stationary points on the PES of the HDA reaction of <b>1b</b> with <b>2b</b> in the <i>endo</i> manner.	122
Figure 4.7	Optimized geometries of the stationary points on the PES of the HDA reaction of <b>1c</b> with <b>2b</b> in the <i>endo</i> manner.	123
Figure 4.8	Optimized geometries of the stationary points on the PES of the HDA reaction of <b>1d</b> with <b>2b</b> in the <i>endo</i> manner.	124
Figure 4.9	Energy profile diagram for the reaction of benzaldehyde with DMB in the presence of different catalysts.	130
Figure 4.10	LUMOs of benzaldehyde complexes with AlCl <sub>3</sub> ( <b>1a</b> ), BF <sub>3</sub> ( <b>1b</b> ), EtAlCl <sub>2</sub> ( <b>1c</b> ) and ZnCl <sub>2</sub> ( <b>1d</b> ) and the HOMOs of DMB ( <b>2a</b> ) and isoprene ( <b>2b</b> ) along with their energies.	133
Figure 4.11	IR spectrum of 3,6-dihydro-4,5-dimethyl-2-(4'-chlorophenyl)-2 <i>H</i> -pyran ( <b>5a</b> )	137
Figure 4.12	<sup>1</sup> H NMR spectrum of 3,6-dihydro-4,5-dimethyl-2-(4'-chlorophenyl)-2 <i>H</i> -pyran ( <b>5a</b> )	138
Figure 4.13	<sup>13</sup> C NMR spectrum of 3,6-dihydro-4,5-dimethyl-2-(4'-chlorophenyl)-2 <i>H</i> -pyran ( <b>5a</b> )	140